

Teagasc National Farm Survey Sustainability Report 2022

OCTOBER 6TH 2023



AGRICULTURAL ECONOMICS AND FARM SURVEY
DEPARTMENT TEAGASC

ISBN: 978-1-84170-691-7



Teagasc National Farm Survey 2022 Sustainability Report

Cathal Buckley and Trevor Donnellan

Agricultural Economics and Farm Surveys Department,

Rural Economy and Development Programme,

Teagasc,

Athenry, Co. Galway,

Ireland

October 6th 2023

ISBN: 978-1-84170-691-7

Contents

Contents	i
Authors and Contact Details	iii
Acknowledgements	iv
Abbreviations	v
Glossary of Terms	vi
What's In the Report	viii
Farm Classifications	ix
List of Tables	x
List of Figures	xi
Executive Summary	xv
Infographics	xvii



Contents

1. Introduction - Agricultural Sustainability	2
2. Measuring Farm Level Sustainability	2
3. Description of the Sustainability Indicators	5
4. Interpretation of the Sustainability Indicators	17
5. Dairy Farm Sustainability 2022	18
6. Cattle Farm Sustainability 2022	28
7. Sheep Farm Sustainability 2022	37
8. Tillage Farm Sustainability 2022	45
9. Farm System Comparisons 2022	52
10. Time Series Comparisons 2015-2022	57
Appendix 1 Individual year results 2017-2022	76



Authors and Contact Details

Cathal Buckley and Trevor Donnellan

The Teagasc National Farm Survey is located in Athenry, Co. Galway, with data recording staff also based at various other Teagasc locations throughout Ireland.



Address

Agricultural Economics and Farm Surveys Department,
Teagasc, Athenry, Co Galway, H65 R718, Ireland



Contact

Tel: +353 91 845 281

Email: nfs@teagasc.ie



Acknowledgements

The authors wish to thank all who contributed to the Teagasc National Farm Survey 2022 - the farmers who participate voluntarily, the Central Statistics Office who select the sample and provide the population weights. Our appreciation is extended to the Teagasc research staff involved in the collection and validation of the farm data: J. Brennan, A. Curley, L. Deane, T. Doyle, P. Harnett, S. Hegarty, G. Kenny, J. Maughan, J. McConnon, K. McNamara, M. Murphy, M. Nicholson, J. Robinson, D. Schilder and J. Teehan as well as Muriel Clarke for the administration of the survey. Best wishes to J. Colgan and P. Healy on their retirement, after illustrious careers with the National Farm Survey team.

Monetary Amounts in Nominal Terms

Monetary figures in this report are presented in nominal terms. This is relevant when considering incomes over time, as inflation, even at a low rate, accumulates over several years and erodes the purchasing power of money. For much of the last decade inflation has been very low in Ireland. However, in 2021 and in 2022, the inflation rate has increased sharply. This is important when considering the change in nominal amounts over recent years.

Interpreting the Box Plots

Some of the data contained in this report are presented in a series of boxplots. These help provide a more in-depth description of the data. In each boxplot, the green shaded boxes are representative of the farms that lie between the 25th and 75th percentile of the NFS farm population. The line within the box represents the median (middle) data point, i.e. half of all farms lie either above or below this point. The tails at either end correspond to the minimum and maximum data points with extreme outliers removed.

Abbreviations

CH ₄	Methane
CO ₂	Carbon dioxide
CO ₂ e	Carbon dioxide equivalent
CSO	Central Statistics Office
ESD	EU Effort Sharing Decision
FPCM	Fat and protein corrected milk
GHG	Greenhouse gases
GM	Gross Margin
GWP	Global Warming Potential
IPCC	Intergovernmental Panel on Climate Change
LCA	Life Cycle Analysis
N	Nitrogen
NH ₃	Ammonia
N ₂ O	Nitrous oxide
NFS	National Farm Survey
NUE	Nitrogen use efficiency
P	Phosphorus
PUE	Phosphorus use efficiency

Glossary of Terms

CO2 equivalent: For reporting purposes all non-carbon dioxide (CO₂) emissions of GHG are converted to CO₂ equivalents using appropriate global warming potentials (GWP100) for CH₄ and N₂O which are respectively 28 and 265 times greater than CO₂.

Direct Costs: Costs directly incurred in the production of a particular enterprise, e.g., fertilisers, seeds and feeding stuffs.

Fat and Protein Corrected Milk (FPCM): This is the functional unit used for carbon foot printing dairy output on farm. It adjusts kg/litres of milk to allow for the level of milk solids produced which is standardized to 4% fat and 3.3% true protein per kilogramme of milk.

Greenhouse Gases (GHG) per farm: The average amount of territorial greenhouse gas emissions (CO₂, N₂O, CH₄) produced in a particular farm type. The approach follows the recognised IPCC methodology used in the calculation on the National GHG inventory

Greenhouse Gases (GHG) per hectare: The average amount of territorial greenhouse gas emissions (CO₂, N₂O, CH₄) produced in a particular farm system expressed on a per hectare basis. The approach follows the recognised IPCC methodology used in the calculation on the National GHG inventory

Greenhouse Gases (GHG) per unit of output: The average amount of territorial greenhouse gas emissions (CO₂, N₂O, CH₄) associated with the production of a specific type of agricultural product, expressed as kg CO₂ equivalent per kg of produce (e.g. per kg beef, milk). The approach follows the recognised IPCC methodology used in the calculation on the National GHG inventory

Gross Output: Gross output for the farm is defined as total sales less purchases of livestock, plus value of farm produce used in the house, plus receipts for hire work, services, fees etc. It also includes net change in inventory, which in the case of cows, cattle and sheep is calculated as the change in numbers valued at closing inventory prices. All non-capital grants, subsidies, premiums, headage payments are included in gross output in this report.

Gross Margin: Gross output minus direct costs.

Glossary of Terms

Global Warming Potential: When counting the emissions of various greenhouse gases they are brought to a common base, or CO₂ equivalent. This common base is arrived at by applying a global warming potential (GWP) to each gas (e.g. N₂O, CH₄). The GWP for CH₄ and N₂O used in this report are those published by the IPCC AR5 report.

Labour Unit: One labour unit is defined as at least 1,800 hours worked on the farm by a person over 18 years of age. Persons under 18 years of age are given the following labour unit equivalents:

16-18 years: 0.75

14-16 years: 0.50

Please note: An individual cannot exceed one labour unit even if he/she works more than 1,800 hours on the farm.

Life Cycle Analysis: An alternative method to the IPCC approach to measuring carbon is the Life-Cycle Assessment approach which accounts for emissions through the entire food production supply chain.

Nitrogen balance: (per hectare farmed), is used as an indicator of the potential magnitude of nitrogen surplus which reflects the risk of nutrient losses to water bodies all other things being equal. It is calculated on the basis of N inputs less N outputs on a per hectare basis at the farm gate level.

Nitrogen use efficiency: is an indicator used to highlight the proportion of N retained in the farm system (N outputs / N inputs). This is a generic measure allowing comparison across disparate farm types at the farm gate level.

Phosphorus balance: (per hectare farmed), is used as an indicator of the potential magnitude of phosphorus surplus which may result in nutrient losses to water bodies all other things being equal. It is calculated on the basis of P inputs less P outputs on a per hectare basis at the farm level.

Phosphorus use efficiency: is used to highlight the proportion of P retained in the farm system (P outputs / P inputs). This is a generic measure allowing comparison across different farm types.



What's in the Report?

Farm Sustainability Performance

- Dairy Farm 2022
- Cattle Farm 2022
- Sheep Farms 2022
- Tillage Farms 2022
- Time series 2015-2022

Farm Categorisation

- Farms typically produce more than one type of agricultural output. In the National Farm Survey farms are categorised into farm types according to their principal output.
- In this Report for 2022, the survey sample is representative of a population of 85,806 farms in Ireland.

Key Performance Indicators

- A broad range of economic, environmental and social indicators is provided, including farm income, labour input, GHG emissions, ammonia emissions, N & P use efficiency, household characteristics and technology adoption.

Farm Classification

Teagasc collects farm data through the National Farm Survey, principally in fulfilment of Ireland's obligation as a member of the European Union. However, the National Farm Survey has evolved over the years to produce a comprehensive list of measures relating to farm sustainability, covering economic, social and environmental performance metrics.

This report focusses mainly on the economic sustainability of Irish agriculture. A dedicated Sustainability Report covering the wider suite of sustainability metrics will be produced later in the year.

The results of the Teagasc National Farm Survey (NFS) can be decomposed in various ways. One of the most common ways in which the results are presented is on a system basis. By system, the NFS farms are categorised into one of six farm types: Dairy, Cattle Rearing, Cattle Other, Sheep, Tillage and Mixed Livestock. Given that individual farms typically have more than one farm enterprise, a rigorous basis for categorising farms into each system is required.

The method of classifying farms into farming systems, is based on the EU farm typology, as set out in Commission Decision 78/463 and its subsequent amendments. The approach is utilised by all members of the EU Farm Accountancy Data Network (FADN).

The methodology assigns a standard output (SO) to each type of animal and each hectare of crop on the farm. Farms are then classified into groups, according to the proportion of total SO which comes from each enterprise. It is important to appreciate that system titles refer to the **dominant** enterprise in each group. For example, the cattle rearing system refers to those farms where the greater proportion of the farm's activity relates to suckler beef production. There are many other farms (including those in the dairy, sheep and tillage systems) that have a cattle enterprise, but where the main enterprise of the farm is not cattle production. Similarly, there will be farms that have sheep, but where cattle is the main enterprise. Tillage farms will sometime also have a secondary enterprise, most often a cattle production system. The mixed nature of many Irish farms is reflected in the individual contribution of livestock and crop categories to farm gross output.

List of Tables

Table 3.1: Overview of Economic Indicators	35
Table 3.2: Overview of Environmental Indicators	37
Table 3.3: Overview of Social indicators	42
Table 3.4: Overview of Innovation indicators	43
Table A 1: Sustainability Indicator results for Dairying Farms 2017-2022	106
Table A 2: Sustainability Indicator results for Cattle Farms 2017-2022	108
Table A 3: Sustainability Indicator results for Sheep Farms 2017-2022.....	110
Table A 4: Sustainability Indicator results for Tillage Farms 2017-2022	112
Table A 5: Sustainability Indicator results for All Farms 2017-2022.....	114

List of Figures

Figure 3-1: Sustainability Overview	34
Figure 3-2: An illustration of some of the major agricultural greenhouse gas emissions.....	38
Figure 4-1: Example Boxplot Gross Margin € per hectare.....	46
Figure 4-2: Example Bar Chart Proportion of farms	46
Figure 5-1: Economic Return and Profitability of Land: Dairy Farms	48
Figure 5-2: Economic Viability: Dairy Farms.....	48
Figure 5-3: Productivity of Labour: Dairy Farms.....	48
Figure 5-4: Market Orientation: Dairy Farms	48
Figure 5-5: Family Farm Income per hectare: Dairy Farms	49
Figure 5-6: Agricultural GHG Emissions for the average Dairy Farm	49
Figure 5-7: Agricultural GHG Emissions per hectare: Dairy Farms	49
Figure 5-8: Agricultural GHG Emissions per kg of Milk: Dairy Farms	49
Figure 5-9: Agricultural GHG Emissions per kg of FPCM: Dairy Farms	50
Figure 5-10: Energy use related GHG	50
Figure 5-11: Energy use related GHG Emissions per kg of Milk: Dairy Farms	50
Figure 5-12: Energy GHG Emissions per kg of FPCM: Dairy Farms.....	50
Figure 5-13: Total LCA based GHG emissions (Agriculture & Energy) per kg of FPCM: Dairy Farms	51
Figure 5-14: Total Ammonia Emissions for the average Dairy Farm	51
Figure 5-15: Ammonia Emissions kg per hectare: Dairy Farms.....	51
Figure 5-16: Ammonia Emissions per kg of FPCM: Dairy Farms.....	51
Figure 5-17: Ammonia Emissions per kg of Milk: Dairy Farms	52
Figure 5-18: N Balance per ha: Dairy Farms	52
Figure 5-19: N Use Efficiency: Dairy Farms.....	52
Figure 5-20: P Balance per ha: Dairy Farms.....	52
Figure 5-21: P Use Efficiency: Dairy Farms	53
Figure 5-22: Household Vulnerability: Dairy	53
Figure 5-23: Agricultural Education: Dairy	53
Figure 5-24: Isolation Risk: Dairy Farms	53
Figure 5-25: High Age Profile: Dairy Farms.....	54
Figure 5-26: Hours Worked on farm: Dairy Farm Operator	54
Figure 5-27: Total Hours Worked: Dairy Farm Operator.....	54
Figure 5-28: Milk Recording: Dairy Farms	54
Figure 5-29: Discussion Group: Dairy Farms	55
Figure 5-30: Spring Slurry: Dairy Farms.....	55
Figure 5-31: Slurry applied by Low emissions slurry spreading methods: Dairy Farms	55
Figure 5-32 Protected Urea Use: Dairy Farms.....	55
Figure 5-33: Liming: Dairy Farms.....	56
Figure 5-34: Reseeding: Dairy Farms.....	56
Figure 6-1: Economic Return and Profitability of Land: Cattle Farms	58
Figure 6-2: Economic Viability: Cattle Farms.....	58
Figure 6-3: Productivity of Labour: Cattle	58
Figure 6-4: Market Orientation: Cattle Farms.....	58
Figure 6-5: Family Farm Income per hectare: Cattle Farms	59

Figure 6-6: Agricultural GHG Emissions for the average Cattle Farm	59
Figure 6-7: Agricultural GHG Emissions per hectare: Cattle Farms.....	59
Figure 6-8: Agricultural GHG Emissions per kg live-weight beef produced: Cattle Farms	60
Figure 6-9: Energy GHG Emissions per hectare: Cattle Farms.....	60
Figure 6-10: Energy use related GHG Emissions per kg live-weight beef: Cattle Farms	60
Figure 6-11: Total Ammonia Emissions for the average Cattle Farm	60
Figure 6-12: Ammonia Emissions per hectare: Cattle Farms	61
Figure 6-13: Ammonia Emissions per kg live-weight beef produced: Cattle Farms.....	61
Figure 6-14: N Balance per ha: Cattle Farms	61
Figure 6-15: N Use Efficiency: Cattle Farms	61
Figure 6-16: P Balance per ha: Cattle Farms	62
Figure 6-17: P Use Efficiency: Cattle Farms	62
Figure 6-18: Household Vulnerability: Cattle	62
Figure 6-19: Agricultural Education: Cattle Farms	62
Figure 6-20: Isolation Risk: Cattle Farms	63
Figure 6-21: High Age Profile: Cattle Farms	63
Figure 6-22: Hours Worked on Farm: Cattle Farm Operator	63
Figure 6-23: Total Hours Worked: Cattle Farm Operator.....	63
Figure 6-24: Spring Slurry Application: Cattle Farms.....	64
Figure 6-25: Low emission slurry spreading: Cattle Farms.....	64
Figure 6-26: Protected Urea use: Cattle Farms	64
Figure 6-27: Liming: Cattle Farms.....	64
Figure 6-28: Discussion Group: Cattle Farms	65
Figure 6-29: Re-seeding: Cattle Farms	65
Figure 7-1: Economic Return and Profitability of Land: Sheep Farms.....	67
Figure 7-2: Economic Viability: Sheep Farms	67
Figure 7-3: Productivity of Labour: Sheep Farms	67
Figure 7-4: Market Orientation: Sheep Farms.....	67
Figure 7-5: Family Farm Income per hectare: Sheep Farms.....	68
Figure 7-6: Agricultural GHG Emissions for the average Sheep Farms	68
Figure 7-7: Agricultural GHG Emissions per hectare: Sheep Farms	68
Figure 7-8: Agricultural GHG Emissions per kg live-weight produced: Sheep Farms.....	68
Figure 7-9: Energy GHG Emissions per hectare: Sheep Farms	69
Figure 7-10: Energy use related GHG Emissions per kg live-weight produced: Sheep Farms.....	69
Figure 7-11: Total Ammonia Emissions for the average Sheep Farm	69
Figure 7-12: Ammonia Emissions per hectare: Sheep Farms	69
Figure 7-13: Ammonia Emissions per kg live-weight produced: Sheep Farms	70
Figure 7-14: N Balance per ha: Sheep Farms.....	70
Figure 7-15: N Use Efficiency: Sheep Farms	70
Figure 7-16: P Balance per ha: Sheep Farms	70
Figure 7-17: P use efficiency: Sheep Farms	71
Figure 7-18: Household Vulnerability: Sheep Farms	71
Figure 7-19: Agricultural Education: Sheep Farms	71
Figure 7-20: Isolation Risk: Sheep Farms.....	71
Figure 7-21: High Age Profile: Sheep Farms	72

Figure 7-22: Hours Worked On Farm: Sheep Farm Operators	72
Figure 7-23: Total Hours Worked: Sheep Farm Operator	72
Figure 7-24: Spring Slurry: Sheep Farms	72
Figure 7-25: Protected Urea use: Sheep Farms.....	73
Figure 7-26: Liming: Sheep Farms	73
Figure 7-27: Reseeding: Sheep Farms	73
Figure 7-28: Discussion Group: Sheep Farms	73
Figure 8-1: Economic Return and Profitability of Land: Tillage Farms	75
Figure 8-2: Economic Viability: Tillage Farms	75
Figure 8-3: Productivity of Labour: Tillage Farms.....	75
Figure 8-4: Market Orientation: Tillage Farms	75
Figure 8-5: Family Farm Income per hectare: Tillage Farms	76
Figure 8-6: Agricultural GHG Emissions for the average Tillage Farm	76
Figure 8-7: Agricultural GHG Emissions per hectare: Tillage Farms.....	76
Figure 8-8: Energy GHG Emissions per hectare: Tillage Farms.....	76
Figure 8-9: Total Ammonia Emissions for the average Tillage Farm	77
Figure 8-10: Total Ammonia Emissions per hectare: Tillage Farms	77
Figure 8-11: N Balance per hectare: Tillage Farms.....	77
Figure 8-12: N Use Efficiency: Tillage Farms.....	77
Figure 8-13: P Balance per hectare: Tillage Farms	78
Figure 8-14: P Use Efficiency: Tillage Farms	78
Figure 8-15: Household Vulnerability: Tillage	78
Figure 8-16: Agricultural Education: Tillage Farms.....	78
Figure 8-17: Isolation Risk: Tillage Farms	79
Figure 8-18: High Age Profile: Tillage Farms.....	79
Figure 8-19: Hours Worked on Farm: Tillage Farms.....	79
Figure 8-20: Total Hours Worked: Tillage Farm Operator.....	79
Figure 8-21: Liming: Tillage Farms.....	80
Figure 8-22: Discussion Group: Tillage Farms	80
Figure 8-23: Break Crops: Tillage	80
Figure 9-1: Economic Sustainability: Farm System Comparison 2022 (average per system)	82
Figure 9-2: Environmental Sustainability: Farm System Comparison 2022 (average per system)	84
Figure 9-3: Social Sustainability: Farm System Comparison 2022 (average per system).....	85
Figure 10-1: Economic Returns to Land: 3 year rolling average 2017-2022	88
Figure 10-2: Profitability of Land: 3 year rolling average 2017-2022.....	88
Figure 10-3: Family Farm income: 3 year rolling average 2017-2022	89
Figure 10-4: Productivity of Labour: 3 year rolling average 2017-2022.....	89
Figure 10-5: Percentage of Output Derived from Market: 3 year rolling average 2017-2022.....	90
Figure 10-6: Economic Viability: 3 year rolling average 2017-2022	90
Figure 10-7: Ag. Greenhouse Gas Emissions per hectare: 3 year rolling average 2017-2022	91
Figure 10-8: Energy Greenhouse Gas Emissions per hectare: 3 year rolling average 2017-2022.....	91
Figure 10-9: Ag. GHG Emissions per Euro output: 3 year rolling average 2017-2022.....	92
Figure 10-10: Energy related GHG Emissions per Euro output: 3 year rolling average 2017-2022	93
Figure 10-11: Kg of Ammonia Emissions per hectare: 3 year rolling average 2017-2022	93
Figure 10-12: Ammonia (NH ₃) Emissions per Euro Output: 3 year rolling average 2017-2022	94

Figure 10-13: Nitrogen Balance per ha: 3 year rolling average 2017-2022	94
Figure 10-14: Phosphorus (P) Balance per ha: 3 year rolling average 2017-2022	95
Figure 10-15: Nitrogen Use Efficiency: 3 year rolling average 2017-2022	95
Figure 10-16: Phosphorus Use Efficiency: 3 year rolling average 2017-2022	96
Figure 10-17: Farm Household Vulnerability: 3 year rolling average 2017-2022	97
Figure 10-18: Isolation Risk: 3 year rolling average 2017-2022 (average per system)	97
Figure 10-19: High Age Profile: 3 year rolling average 2017-2022 (average per system)	98
Figure 10-20: Hours Worked On Farm Per Annum: 3 year rolling average 2017-2022 (average per system) .	98
Figure 10-21: Total Hours Worked Per Annum: 3 year rolling average 2017-2022 (average per system)	99
Figure 10-22: Formal Agricultural Education: 3 year rolling average 2017-2022 (average per system)	99
Figure 10-23: Ag. GHG Emissions per kg FPCM: 2017-2022 Dairy Farms.....	100
Figure 10-24: Ag. GHG Emissions per kg live-weight beef produced: 2017-2022 (Cattle Farms)	100
Figure 10-25: Ag. GHG Emissions per kg live-weight sheep produced: 2017-2022 Sheep Farms.....	100
Figure 10-26: Energy use related GHG emissions per kg FPCM: 2017-2022 Dairy Farms.....	100
Figure 10-27: Energy use related GHG emissions per kg live-weight beef produced: 2017-2022 Cattle Farms	101
Figure 10-28: Energy use related GHG emissions per kg live-weight sheep produced: 2017-2022 Sheep Farms	101
Figure 10-29: Ammonia emissions per kg FPCM: 2017-20223 year rolling average Dairy Farms.....	101
Figure 10-30: Ammonia emissions per kg live-weight beef produced: 2017-2022 Cattle Farms	101
Figure 10-31: Ammonia emissions per kg live-weight sheep produced: 2017-2022Sheep Farms	101

Executive Summary

This report provides the latest available information on the sustainability performance of farms in Ireland, based on detailed analysis of data collected through the Teagasc National Farm Survey. Economic, Social, Environmental and Innovation sustainability metrics are produced for Dairy, Cattle, Sheep and Tillage farms in 2022. The report also includes time series results over several years, which allows an assessment of how farm sustainability has changed temporally.

Economic sustainability

- Consistent with the established trend of earlier years, Dairy farms remain an economic powerhouse in Irish agriculture. Average **economic returns per hectare** in Dairy tend to be multiples of those in the other farm systems in Ireland and the gap between Dairy and the other farm systems has tended to widen since the EU decided to eliminate the milk quota system. The report indicates that this gap widened further in 2022, which saw a significant spike in milk prices, due to the slow growth in milk production globally, allied with strong international dairy demand.
- Varying amount of labour are required across different farm systems. Expressing farm incomes on a unit of family labour basis addresses this issue. On making this adjustment Dairy and Tillage farms can be considered as relatively comparable in income terms. The results show that both of these farm system types considerably outperform the drystock farm systems in economic terms.

Social sustainability

- Again reflecting established trends, Dairy continues to exhibit a stronger performance in terms of **social sustainability** relative to other farm systems. Dairy tends to be associated with a lower **isolation risk** (living alone). Fewer dairy farm households have a **high age profile** in comparison with other farm systems. Tillage farms also tend to generally outperform livestock farms on these social sustainability metrics.
- However, dairy farming is typically a very labour intensive system. In terms of **labour input**, on average the main dairy farm operator works significantly more **hours per year** than the average farm operator in the other farm systems. Given the hours required on farm, relatively few dairy farmers work off farm. However, even when the labour from time spent working off farm (which can be significant for drystock systems) is combined with time spent working on-farm, the labour input of dairy farm operators tends to exceed that of farm operators of all other farm systems.

Environmental sustainability

a) Greenhouse gas emissions

- **Dairy:** Even though herd sizes increased in 2022, **total farm and per hectare GHG emissions** on the average dairy farm declined, largely due to a significant decrease in chemical N fertiliser use. The **GHG emissions intensity of milk production** (CO₂ equivalent per kilogramme of Fat and Protein Corrected Milk) also improved. Effectively this means that the average kilogramme of milk on Irish dairy farms was produced with a lower carbon footprint in 2022.
- **Non-Dairy Systems: Farm level and per hectare level GHG emissions** on cattle, sheep and tillage farms also declined in 2022 on the back of reduced chemical N fertiliser use. In percentage terms, the reduction in N fertiliser use was larger on drystock farms than on dairy farms, but in absolute terms, the reduction on dairy farms was larger. This reflects the fact that the average dairy farms typically use close to 3 times as much chemical N per hectare compared to the average drystock farm. The decline in chemical N use was most likely driven by the high N fertiliser prices, which emerged in the second half of 2021 and persisted throughout 2022. However, the advice to farmers now strongly emphasises reduced fertiliser use on both economic and environmental grounds, which may mean that the observed reduction in fertiliser use in 2022 could well be maintained.

b) Ammonia emissions

- On dairy farms ammonia emissions per farm and per hectare increased in 2022 relative to 2021, but the level in 2022 was below the longer terms trend of years before 2021. The increase was driven by the **composition of N fertiliser** used, with a move away from the use of CAN towards straight urea, which is a much bigger emitter of ammonia. Ammonia emissions on other farm system tended to decline in 2022 on due to reduced chemical N use and increased adoption of technologies like **low emission slurry spreading** (LESS). In spite of the increased usage of LESS on drystock farms, the use of that technology is still less prevalent than on dairy farms, indicating that there is room for further progress in this area on drystock farms.

c) Nitrogen balance and use efficiency

- **Across all farm systems, N surpluses** declined and **N use efficiency** improved in 2022. This improvement was achieved through reduced chemical N use and increased best practice technology adoption. These metrics tend to be significantly influenced by variability in weather conditions from one year to the next, although improved N management on farms also plays a role.


Innovation

- There was an increase in the percentage of dairy and cattle farms applying lime in 2022. **Lime application** is important in optimising the soil's pH level, which affects the availability of nutrients in the soil and can lead to a reduced requirement for fertiliser use.
- There was also a significant transition towards the **use of LESS equipment** for slurry application. In 2022, 75% of slurry on the average dairy farm and 34% on the slurry on the average cattle farm was applied via LESS.
- However, while the percentage of chemical N applied in the form of **protected urea** is growing on dairy and cattle farms, it remains at a relatively low level in absolute terms.



Gross Output per ha 2022

€6,005




Gross Margin per ha 2022

€3,509




Family Farm Income per ha 2022

€2,332




Productivity of Labour 2022

€109,003



Market Orientation 2022

94%



Viability 2022

93%




Source: Teagasc National Farm Survey



Household Vulnerability 2022

4%




Isolation 2022

6%




High Age Profile 2022

18%



Hours Worked on Farm 2022

2,498




Total Hours Worked 2022

2,657



Agricultural Education 2022

83%



Source: Teagasc National Farm Survey




CO2 eq per ha 2022

9.4




CO2 Eq per kg FPCM 2022

0.82




CO2 Eq per euro of output 2022

1.7



NH3 kg per ha 2022

46.9



N Balance kg per ha 2022

158.6




Source: Teagasc National Farm Survey



Gross Output per ha 2022

€1,740



Gross Margin per ha 2022

€845




Family Farm Income per ha 2022

€372



Productivity of Labour 2022

€17,154



Market Orientation 2022

70%



Viability 2022

27%



Source: Teagasc National Farm Survey



Household Vulnerability 2022

33%




Isolation 2022

17%




High Age Profile 2022

41%



Hours Worked on Farm 2022

1,370




Total Hours Worked 2022

2,139



Agricultural Education 2022

49%



Source: Teagasc National Farm Survey



CO2 eq per ha 2022

4.4




CO2 eq per kg of liveweight 2022

9.4




CO2 per euro of output 2022

3.5




NH3 kg per ha 2022

22.0



N Balance kg per ha 2022

43.6



Source: Teagasc National Farm Survey



Gross Output per ha 2022

€1,475



Gross Margin per ha 2022

€625




Family Farm Income per ha 2022

€340



Productivity of Labour 2022

€14,890



Market Orientation 2022

65%



Viability 2022

26%




Source: Teagasc National Farm Survey



Household Vulnerability 2022

29%




Isolation 2022

13%




High Age Profile 2022

28%



Hours Worked on Farm 2022

1,477




Total Hours Worked 2022

2,246



Agricultural Education 2022

61%



Source: Teagasc National Farm Survey



CO2 eq per ha 2022

3.3



CO2 per kg liveweight 2022

7.8




CO2 per euro of output 2022

3.5



NH3 kg per ha 2022

12.7



N Balance kg per ha 2022

30.4



Source: Teagasc National Farm Survey



Gross Output per ha 2022

€2,812




Gross Margin per ha 2022

€1,581




Family Farm Income per ha 2022

€1,043




Productivity of Labour Unit 2022

€97,910



Market Orientation 2022

82%



Viability 2022

79%



Source: Teagasc National Farm Survey



Household Vulnerability 2022

10%




Isolation 2022

18%



High Age Profile 2022

25%



Hours Worked on Farm 2022

1,416




Total Hours Worked 2022

2,133



Agricultural Education 2022

63%



Source: Teagasc National Farm Survey




CO2 eq per ha 2022

1.9




CO2 eq per euro of output 2022

0.9



NH3 kg per ha 2022

9.1



N Balance kg per ha 2022

40.5



P Balance kg per ha 2022


3.6




Source: Teagasc National Farm Survey

Social Sustainability Special Focus 2022


When did you last take a break from the farm?




When did you last visit a doctor?



Daily contact with people outside the farm?




Provide/receive help on farm




Farm Labour

Labour hours and Farm size



Proportion of farms with female labour input 2022



These social sustainability metrics for 2022 are based on data from a special survey of a sample of 611 farms, representing over 85,000 farms nationally.

Farm Sustainability 2022

Detailed Results

1 Introduction - Agricultural Sustainability

Civilization faces a grand challenge in trying to feed a growing human population, while minimising the environmental impacts of food production, especially in the context of climate change, deteriorating water quality and biodiversity loss. To feed a growing global population in a sustainable way, agricultural output must increase without reducing the capacity for future production or compromising the environment. This is the overarching objective of the EU Farm to Fork Strategy published in May 2020 (European Commission, 2020). Separately, Ireland has passed legislation (Climate Action and Low Carbon Development Act, 2021) which sets down a target of a 51% reduction in national greenhouse gas emissions by 2030, with sectoral ceiling allocated by Government (Government of Ireland, 2022) to the Agricultural sector that are equivalent to a target of a 25% GHG emissions reduction from the sector by 2030.

Agricultural systems are complex and tend to have multiple objectives and wide-reaching effects, which must be considered holistically. To measure and track the diverse elements of Irish farm systems, this report considers Irish **agricultural sustainability** (and its component farm systems) in terms of its **economic, environmental, social** and **innovation** dimensions. The report recognises that the multi-dimensional nature of sustainability means that the exclusive pursuit of improvements in any one of these dimensions alone, is unlikely to deliver a sustainable outcome.

2 Measuring Farm Level Sustainability

The measurement of agricultural sustainability is challenging, as it is a broad concept covering diverse elements, which may vary through time and space. In geographic terms Ireland is small. However, its low human population density and temperate maritime climate means that it has a comparatively large agricultural sector which, given the climate, is focused on grassland agriculture, supporting dairy, cattle and sheep production in particular. Due to climate and topography, Ireland's arable sector is small by comparison with that found in other EU member states.

However, these grassland systems, are heterogeneous, with substantial variations between the typical farms in each farm system in terms of farm size, stocking rates and input usage. Relevant indicators which capture this diversity are required to assess the sustainability status of Irish farms through time. Such metrics can highlight particular areas of concern or trends through time and indicate areas where improvement may be needed. The need to ensure that robust, comprehensive multi-dimensional sustainability data are available for Irish agriculture was recognised over a decade ago. Ireland is fortunate to be at the forefront in Europe in the development and use of wide ranging sustainability metrics for agriculture. Many EU Member States, including some of those with the largest agricultural sectors, have yet to make significant progress in sustainability data provision. The challenge these Member States face is that the challenge of sustainability data collection and provision is considerable and urgent progress is required.

Deriving and maintaining a sustainability indicator set is difficult, as it requires detailed, accurate and consistent farm-level measurements and data across a wide range of physical, socioeconomic and demographic farm attributes. The Teagasc National Farm Survey (NFS) has evolved in response to the changing needs of stakeholders to provide such a dataset. The NFS is a nationally representative sample of approximately 840 farms from across Ireland. Data from the Teagasc NFS represent the Irish component of the European Union's Farm Accountancy Data Network (FADN) dataset¹. However, the data collected in the Teagasc NFS surpasses the requirements of FADN, giving the Teagasc NFS dataset much more capacity to measure and track developments in agricultural sustainability. The Teagasc NFS collects data on an ongoing basis, with the results published annually. A weighting system, based on data published by the Central Statistics Office (CSO), reflective of the national farm population is applied to the data from the individual NFS farms. In this way, national level representation is achieved in terms of size and farm type for the principal farm systems in Ireland. This population weighting is important to ensure that meaningful and accurate aggregations of farm types are made at an appropriate scale (for example, based on farm system type). It also means that the survey results are capable of accurately highlighting synergies and/or trade-offs between different indicators, depending on how farms are managed.

Within the Teagasc NFS, farms are classified into major farming systems according to the standardised EU typology as set down by European Commission regulations and applied by the EU FADN (a more detailed explanation and the correspondence between the farm systems used in the NFS and the farm types set out in the EU farm typology can be found in the Teagasc National Farm Survey Report (Donnellan et al., 2020)). This report presents results for the four dominant land based farm systems in Ireland, namely, dairy, cattle, sheep and tillage.

As the required data are produced on an annual basis, it is possible to generate and compare indicators over time. As methodologies are updated and data requirements evolve to reflect scientific developments, the entire time series of sustainability metrics are revised to reflect current scientific knowledge. This is evident in the time-series analysis for key indicators presented in the report, which are revised to reflect recognised developments in the measurement of sustainability. It is expected that, based on scientific advances and emerging areas of interest (e.g. in both a scientific and policy context), the sustainability indicator set will continue to evolve to maximise its relevance. The aim is that as indicator methodologies develop, they will still be capable of being generated using Teagasc NFS data, ensuring an on-going accurate inter-temporal assessment of the sustainability performance of Irish farm systems.

Furthermore, as the NFS is part of the EU FADN, there is scope for comparative analysis with the sustainability performance of farms in other EU Member States. Indeed, the EU Farm to Fork strategy (EU Commission, 2020) proposes to develop the EU FADN into a EU Farm Sustainability Data Network (FSDN), with a view to collecting data on sustainability indicators and reporting these in a common framework across the EU. EU legislation to bring the FSDN into existence has been drafted in 2022 and is being finalised in 2023. Preparations for the

¹ The Teagasc NFS sampling frame is restricted to farms over €8,000 of standard output (equivalent to 6 dairy cows, 6 hectares of wheat or 14 suckler cows). A total of circa 85,000 farms are represented in this study for 2021.

transition to FSDN have begun across the EU and the Teagasc NFS is leading the way on this, as evidenced by the content of this report.

3 Description of Sustainability Indicators

The indicators described here follow on those published in previous Teagasc sustainability reports (Hennessy et al., 2013; Lynch et al., 2016; Buckley et al., 2019; Buckley & Donnellan, 2020a; 2020b; 2021, 2022). Updates presented here reflect methodological refinements, as well as additional data on agricultural activities on Irish farms collected and published by the Teagasc NFS. For instance, some GHG emission factors have been updated leading to a slight adjustment to GHG results across farm systems relative to previously published values for past years. Hence, the historical time series for some of the sustainability indicators presented in the current report differ and supersedes those presented in earlier Teagasc Sustainability reports (Buckley et al., 2019; Buckley & Donnellan, 2020a; 2020b, 2021, 2022). This approach to revising historic sustainability indicators is to ensure they fully reflect our current scientific knowledge. It mirrors the approach used by Ireland’s Environmental Protection Agency (EPA) in their national inventory reporting for Ireland and is hence consistent with international best practice.

As depicted in Figure 1 and described in the following section, the Teagasc Sustainability Report’s indicators are grouped into four categories: **economic, environmental, social and innovation.**

Figure 3-1: Sustainability Overview



3.1 Economic Indicators

Economic viability is essential to ensure that a farm system can sustain itself and that farming families are compensated adequately for owned capital and labour employed. At a national level, agriculture is an important component of the Irish economy, particularly in more rural areas. The Teagasc NFS is well equipped to generate economic indicators. Its origins derive from the need to submit data on economic farm performance to the European Commission through the EU FADN, primarily to aid in the understanding of the effectiveness of the Common Agricultural Policy (CAP). The economic sustainability indicator set is, therefore, relatively comprehensive and (largely unconstrained by issues relating to data availability) designed to cover a range of important economic measures. The following economic indicators are presented in the report:

Table 3.1: Overview of Economic Indicators

<i>Indicator</i>			<i>Measure</i>	<i>Unit</i>
Market output	based	gross	Gross output per hectare	€ / hectare
Market margin	based	gross	Market based gross margin per hectare	€ / hectare
Productivity of Labour			Family Farm Income per unpaid labour unit	€ / unpaid labour unit
Economic Viability			Economic viability of farm business	1=viable, 0=not viable
Market Orientation			Output derived from market rather than subsidies	%
Family Farm Income			Family Farm Income per hectare	€ / hectare

a) Market based gross output

This is measured as **gross output (€) per hectare of utilised agricultural area (UAA)**. Gross output is defined as total sales less purchases of livestock, livestock based products & crops, plus the value of farm produce used in the household plus receipts for hire work, service fees etc. It also includes the value of net changes in inventories, which for cows, cattle and sheep are calculated as the change in numbers year on year valued at closing inventory prices. All non-capital grants, subsidies and premium payments are also included in gross output, as are income from land and quota lettings. Inter-enterprise transfers are then deducted in order to avoid double counting of activity. A limitation of this measure is that by definition it can change due to output price inflation. It measures change which is due commodity price volatility, in addition to increased physical output. Therefore care is required in interpreting inter annual changes in this metric.

b) Profitability of Agricultural Land Use

The **market based gross margin** (gross margin excluding grants and subsidies), is where gross margin is defined as gross output less direct costs per hectare. Given that it is profit based, this measure is less susceptible to inflation, since inflation on both the output and input side tend to cancel each other out.

c) Productivity of Family Labour

In the NFS, a distinction is made between the labour of farm family members, which is generally unpaid and therefore is not classified as a production cost, and hired labour, which in accounting terms does represents a production cost to the family farm. The return to unpaid family labour is measured as **family farm income per unpaid family labour unit**. For

consistency in measurement of farm labour input across the EU, one labour unit is defined as a person over 18 years old, working at least 1,800 hours a year (to maintain international consistency in the measure, it is not possible to report in excess of one labour unit per person, even where an individual works more than 1,800 hours). Relative to a labour unit for an adult, labour unit equivalents of 0.75 and 0.5 are used for individuals aged 16-18 years and 14-16 years respectively.

d) Economic Viability

The **economic viability** of a farm business is measured using a **binary variable**. A farm is defined as viable if family labour is remunerated at greater than or equal to the minimum wage, as set down in the under the National Minimum Wage Act, 2000 (as outlined by Government of Ireland, 2021a) and there is also sufficient income generated by the farm to provide an additional five per cent return on non-land based assets employed on the farm.

e) Market Orientation

The market orientation is measured as the **proportion of gross output (€) that is derived from the market** (generally the sales value of the farm's outputs which can referred to as **market based gross output**), as opposed to grants and subsidies, which are treated as a non-market based gross output of the farm. Over time this measure is also sensitive to output price inflation, given that subsidies are more typically a fixed value over time.

f) Family Farm Income

Family Farm Income (FFI) is the return from farming for farm family labour, land and capital. It is a function of gross output plus subsidies less total net expenses. FFI is measured in nominal terms. This needs to be taken into account, when assessing developments in FFI over time, particularly in periods of high inflation such as occurred in 2022.

3.2 Environmental Indicators

Agriculture can generate positive or negative environmental impacts depending on the specific activities undertaken on the farm. Agriculture is the principal land use in Ireland covering 69% of the State's land area (CSO, 2022). Hence, the environmental sustainability of agriculture is key to achieving national level objectives relating to the environment. The current set of NFS based environmental indicators focus on **greenhouse gas (GHG) emissions, ammonia emissions, nitrogen and phosphorus use**. Indicators that are currently under development include, metrics relating to **biodiversity**. Measures of biodiversity, particularly those designed to accurately pick on small changes through time at the individual farm level, are technically challenging to implement and require considerable resources. A measure of biodiversity quantity will be included in future Teagasc sustainability reports once the relevant scientific work needed to establish indicators and consistently collect the related data has concluded.

Table 3.2: Overview of Environmental Indicators

<i>Indicator</i>	<i>Measure</i>	<i>Unit</i>
Ag. GHG emissions per farm	Absolute Ag. based GHG emissions per farm	Tonnes CO ₂ equivalent / farm
Ag. GHG emissions per hectare	Absolute Ag. based GHG emissions per hectare	Tonnes CO ₂ equivalent / hectare
Ag. GHG emissions per kg of output	Ag. based GHG emissions efficiency	kg CO ₂ equivalent / kg output AND kg CO ₂ e / € output
Energy GHG emissions per farm	Absolute energy GHG emissions per farm	Tonnes CO ₂ equivalent / farm
Energy GHG emissions per hectare	Absolute energy GHG emissions per hectare	Tonnes CO ₂ equivalent / hectare
Energy emissions per kg of output	Energy GHG emissions efficiency	kg CO ₂ equivalent / kg output AND kg CO ₂ e / € output
NH₃ emissions per farm	Absolute NH ₃ emissions per farm	Tonnes NH ₃ equivalent / farm
NH₃ emissions per hectare	Absolute NH ₃ emissions per hectare	Tonnes NH ₃ equivalent / hectare
NH₃ emissions per kg of output	NH ₃ emissions efficiency	kg NH ₃ equivalent / kg output AND kg NH ₃ / € output
N balance	N transfer risk	kg N surplus / ha
N use efficiency	N retention efficiency	% N outputs / N inputs
P balance	P transfer risk	kg P surplus / ha
P use efficiency	P retention efficiency	% P outputs / P inputs

3.2.1 Greenhouse gas emissions

To minimise the extent and impact of climate change, action is required to reduce global greenhouse gas emissions. Agriculture is the largest contributor to Irish greenhouse gas emissions by sector, with 38.4% of the national emissions total in 2022 (EPA, 2023a). The agricultural sector must reduce its emissions in the context of Ireland's commitment to reduce national GHG emissions. The Climate Action and Low Carbon Development (Amendment) Act 2021 (Government of Ireland, 2021b) sets out an ambition for a climate neutral economy by 2050 for the state. Agriculture now has a sectoral target of 25% reduction by 2030 (Government of Ireland, 2022).

Two dominant measures of greenhouse gas emissions exist - the IPCC methodology and the LCA methodology. They differ in important ways and definitions of both are provided below.


1. Intergovernmental Panel on Climate Change (IPCC) Methodology: The GHG emissions indicators in this report are in the first instance calculated following the IPCC methodology accounting conventions and Irish emission factors as employed in the 2021 National Inventory Report for Ireland (Duffy et al., 2023). The IPCC based methodology reflects so called territorial emissions by IPCC sector. This means that the IPCC based approach measures the agriculture based emissions that occur on farms in Ireland. This definition differs from the more holistic approach that is typically used in the Life Cycle Assessment approach (which is explained below), which tries to capture emissions in a production process, including emissions associated with inputs to the process from outside the territory.

The three main agricultural GHG emissions categories are **methane (CH₄)** emissions from **enteric fermentation** by ruminant livestock, CH₄ and **nitrous oxide (N₂O)** emissions from the **production and storage of livestock manures**, and N₂O emissions resulting from

the **crop residues, application of manures and synthetic fertilisers** to agricultural soils. **Carbon dioxide (CO₂)** emissions associated with **liming** and **urea application** are also included in the analysis presented in this report.

A complicating factor inherent in an individual **farm based** approach to emissions measurement, (as opposed to a **national aggregate** emissions inventory approach), is that some animals move between farms via inter-farm sales as part of the agricultural production process within a calendar year. Unless this is factored into the calculations, it could lead to an over estimation or underestimation of activity and associated emissions on individual farms. Accordingly, a farm level animal inventory approach is used here, whereby the CH₄ emissions and manure production of each livestock category are adjusted to reflect the portion of the year an animal is present on a particular farm. For reporting purposes, all non-carbon dioxide (CO₂) emissions are converted to **CO₂ equivalents (CO₂e)** using appropriate global warming potentials (GWP) for CH₄ and N₂O which are respectively 28 and 265 times greater than the GWP of CO₂.

Figure 3-2: An illustration of some of the major agricultural greenhouse gas emissions



Emissions resulting from on-farm fuel and electricity use are considered independently of the IPCC’s agricultural emissions category, as they are recognised under a separate IPCC category (Energy). Energy emissions (CO₂ only) are estimated from expenditure on electricity, natural gas and fuels (relevant quantities used are estimated by using national average prices (CSO, 2023a)) and by applying national level emissions factors to these quantities.

Using the IPCC methodology, the main indicators developed include:

- a. Total agricultural emissions per farm:** with emissions calculated for each farm system. These are also disaggregated to show the emissions originating from different farm enterprises (dairy, cattle, sheep and crops).
- b. Agricultural greenhouse gas emissions per unit of output:** derived so that the total emissions of the farm can be decomposed into components relating to each of the farm’s main agricultural outputs (milk, cattle or sheep live-weight and crop outputs).
- c. Agricultural greenhouse gas emissions per hectare & € of output:** In addition, agricultural based GHG emissions per € of output and per hectare are used to illustrate GHG emissions that are generated on farms with dissimilar levels of agricultural output.
- d. Emissions from on-farm energy use per unit of relevant output:** measures emissions from electricity, gas and fuel use associated with agricultural production

activities on the farm. As per the IPCC methodology, these GHG emissions are considered separately from agricultural GHG emissions.

2. LCA Methodology: An alternative method to the **IPCC approach** of measuring GHG emissions is the **Life Cycle Assessment (LCA) approach**, which accounts for emissions through the entire food production supply chain. LCA is a holistic systems approach that aims to quantify the potential environmental impacts, e.g. GHG emissions, generated throughout a product's life cycle, from raw-material acquisition through production, use, recycling and final disposal. The LCA used here accounts for all agricultural GHG emissions from the farm up to when it leaves the farm. It is generally expressed per unit of product produced. The LCA approach attempts to capture all emissions associated with a product. It therefore ignores national boundaries and seeks to enumerate all emissions along the chain, irrespective of country of origin.

Relative to the territorial IPCC approach, considerably more data are required to conduct an LCA study or to produce a carbon footprint analysis for each product produced on a farm. At present such detailed data are only available for dairy farms participating in the NFS and it was only possible to conduct a carbon LCA based footprint analysis of milk production using NFS data. The Teagasc Dairy LCA model was used for this analysis (O'Brien et al., 2014; Herron et al., 2021). This model, which is accredited by the National Carbon Trust (UK) underpins the carbon footprint results from the Bord Bia Sustainable Dairy Assurance Scheme (SDAS). In LCA analysis system boundaries have to be defined to determine what will and will not be included as part of the calculation. The system boundaries of this LCA model are defined to include all emissions associated with the dairy production system up to the point where milk is sold from the farm. The advantage of applying the Dairy LCA model using NFS data is that the Teagasc NFS is nationally representative of Irish milk production and thus reflects the full spectrum of dairy farming conditions in the country and as such allows for the production of a nationally representative LCA based carbon footprint measure.

As with the other indicators presented in this report, emphasis should not be placed on the absolute level of the carbon footprint measure, since this will be the subject of ongoing revision. Of greater relevance is the direction in which the indicator evolves over time. The main objective of this research is to establish indicators through which changes in sustainability performance can be documented and evaluated.

3.2.2 Ammonia

Ammonia (NH₃) is an air pollutant contributing to **eutrophication** and **acidification** of terrestrial and aquatic ecosystems. It is also an **indirect source** of a potent greenhouse gas **nitrous oxide** (Sutton et al., 1992). The EU and its Member States are parties to the Convention on Long-Range Transboundary Air Pollution (CLRTAP), which regulates trans-boundary air pollutants, including NH₃. Within the EU, NH₃ emissions are regulated through the National Emissions Ceiling (NEC) Directive (EU, Commission 2016). Over 99.4% of Ireland's NH₃ emissions originate within agriculture, principally from animal waste and the application of synthetic fertilisers (EPA, 2022). The fact that NH₃ emissions in Ireland come almost exclusively from agriculture means that any future national ammonia reduction target for Ireland would *de facto* represent a reduction target to be achieved by the agriculture sector. From 2020, Ireland has an NH₃ ceiling of 112.2 kilotonnes per annum, representing a 1% NH₃

reduction relative to the 2005 level. A further reduction target of 5% relative to the 2005 level (to a ceiling of 107.6 kilotonnes per annum) is to be achieved by 2030. The national inventory accounting methodology as applied by Ireland's EPA (Duffy et al., 2021), in conjunction with activity data from the NFS, is used to estimate NH₃ emission indicators across different farm systems in this report. The main indicators developed include:

- a. **Total agricultural ammonia emissions per farm:** with emissions calculated for each farm system. These are also disaggregated to show the emissions originating from different farm enterprises (dairy, cattle, sheep and tillage).
- b. **Ammonia emissions per unit of output/hectare:** derived so that the total NH₃ emissions of the farm can be decomposed into components relating to each of the farm's main agricultural outputs (milk, cattle or sheep live-weight and crop outputs). In addition, NH₃ emissions per € of output and per hectare are used to illustrate emissions that are generated on farms with dissimilar levels of agricultural output.

3.2.3 Nutrient Use Efficiency

Nitrogen (N) and **Phosphorus (P)** use indicators follow a **nutrient accounting approach** based on Buckley et al. (2015). N and P **exports** from the farm are **subtracted from imports** to the farm to give a **farm gate balance**. Exports comprise the N and P component of the farms output, which would include milk, crops, wool, manures exported and livestock sold (including livestock for slaughter). Imports are comprised of fertilisers applied, feeds purchased, livestock brought onto the farm and imported organic manures. It should be noted that the N and P indicators do not provide estimates of losses to water, as such losses are complex and driven by site specific biophysical factors and weather conditions. N and P balances are used as an indicator of potential risk of loss of nutrients, all other things being equal, and cover most of the key management decisions over which the farmer has direct control.

Nitrogen use - Nitrogen (N) is an important element in agricultural production, but the loss of excess N poses a significant risk to the aquatic environment. The nitrogen use indicators follow an input-output accounting methodology as described below.

- c. **Nitrogen balance** (per hectare farmed), is used as an indicator of the potential magnitude of nitrogen surplus, which reflects the risk of nutrient losses to water bodies, all other things being equal. It is calculated on the basis of N inputs less N outputs on a per hectare basis at the farm gate level.
- d. **Nitrogen use efficiency (NUE)** is used to highlight the proportion of N retained in the farm system (N outputs / N inputs). This is a generic measure allowing temporal comparisons at the farm gate level.

Phosphorus use - Similar to nitrogen, phosphorus (P) is an important element in agricultural production and its loss poses a significant risk to the aquatic environment. Phosphorus use indicators, like N use indicators, also follow the input-output accounting methodology described previously. However, it should be noted that unlike N, phosphorus can remain in the soils for significant periods of time and is available to be stored and mined, hence P balance and efficiency should be interpreted with caution in the absence of knowledge of the soil P status of the farm.

- a. **Phosphorus balance** (per hectare farmed), is used as an indicator of the potential magnitude of phosphorus surplus which may result in nutrient losses to water bodies all other things being equal. It is calculated on the basis of P inputs less P outputs on a per hectare basis at the farm level.
- b. **Phosphorus use efficiency** (PUE) is used to highlight the proportion of P retained in the farm system ($P \text{ outputs} / P \text{ inputs}$). This is a generic measure allowing temporal comparisons at the farm gate level.

3.3 Social Indicators

A farm will only be sustainable if employment in agriculture can provide a suitable economic return for the labour employed, but also if farm operators and families have an **acceptable quality of life** from their farming and non-farming activities. If farming is not **socially sustainable**, individuals may exit the sector, or there may be a lack of new entrants to farming, with fewer younger people willing to take over farms when older farmers retire from farming. In addition, as agriculture is often the predominant economic activity in many rural areas, the social impacts of a viable farming sector are also important in **maintaining employment and social well-being** in the broader rural community. The design of social sustainability indicators is subjective in nature and further work is ongoing to improve the farmer, animal and community well-being aspects of social sustainability measurement within the Teagasc NFS. Based on the data currently available from the Teagasc NFS, the following indicators are reported:

Table 3.3: Overview of Social indicators

<i>Indicator</i>	<i>Measure</i>	<i>Unit</i>
Household vulnerability	Farm business is not viable and no off-farm employment in the household	Binary variable: 1= vulnerable
Agricultural education	Formal agricultural training received by the farmer	Binary variable, 1= agricultural training received
Isolation Risk	Farmer lives alone	Binary variable, 1=isolated
High Age Profile	Farmer is over 60 years old, and no members of household under 45	Binary variable: 1=high age
Hours worked on-farm	Farm work load of farmer	Hours worked
Total hours worked	Total farm and off-farm work load of the farmer	Hours worked

a) Household vulnerability

The household vulnerability indicator is a **binary indicator**, where a farm is defined as vulnerable if the farm business is **not economically viable** (using the economic viability indicator described earlier), and the farmer or farmer's spouse has **no off-farm employment income source**.

b) Formal agricultural education

This is a **binary indicator** which measures whether or not the farmer has received any **formal agricultural training**, at any level. Agricultural education can be an important factor in farm succession, as well as having a role in influencing wider farm management decisions that can affect other dimensions of farm sustainability (e.g. willingness to adopt new technologies).

c) High Age Profile

Farm households are defined as having a high age profile if the **farmer is aged over 60**, and there are **no members** of the **farm household younger than 45**. This indicator shows whether the farm household is likely to be demographically viable.

d) Isolation risk

Isolation risk is also measured using a **binary variable**, depending on whether or not the **farmer lives alone**.

e) Hours worked on farm

This indicator is the **number of hours worked** by the farmer **on the farm**. It should be noted that this **does not include** time spent in **off-farm employment**.

f) Total Hours worked

This indicator is the **number of hours worked** by the farmer **on and off the farm**. This **includes** hours worked in **off-farm employment**.

3.4 Innovation Indicators

More efficient production has the potential to increase profitability, while reducing negative environmental and social effects, thereby assisting progress towards more sustainable agriculture. Innovations that can lead to increased sustainability may be **novel technologies**, newly developed or applied, or may arise from the **adoption of** established and newly developed **management techniques**. Hence, it is important to measure uptake of such innovations to measure and evaluate whether evolving science and knowledge is being translated into actual farmer practices and secondly that the use of these technologies gives the anticipated environmental, economic or social benefits. As a result, the innovation indicators selected here are a combination of **specific technologies or practices employed** by the farmer, and also reflect **farmer membership in groups** which may be positively associated with increased adoption of broader innovations. The majority of the innovation indicators are scored as **binary variables**, either where a specific technology or practice is used or where a farmer is a member of the given group. Innovation indicators can be especially useful when evaluated in conjunction with those relating to economic or environmental performance, as they will highlight the benefits of specific technologies or behaviours.

Table 3.4: Overview of Innovation indicators

<i>Dairy</i>	<i>Cattle</i>	<i>Sheep</i>	<i>Tillage</i>
Discussion Group Membership	Discussion Group Membership	Discussion Group Membership	Discussion Group Membership
Liming	Liming	Liming	Liming
Spring slurry spreading*	Spring slurry spreading*	Spring slurry spreading*	Break Crop
Protected urea use	Protected urea use	Protected urea use	
Reseeding	Reseeding	Reseeding	
Low emission slurry spreading	Low emission slurry spreading		
Milk Recording			

*(>50% slurry spread during the period January - April)

Dairy innovation indicators

- **Discussion group membership** was selected as indicating the degree of interaction farmers have with farm extension services and their peers. This is reported in binary (yes/no) format.

- **Liming and Reseeding** were identified as important farming practices in grassland management. This is reported in binary (yes/no) format.
- **Spring slurry spreading** (spreading at least 50% of total slurry between January and April) was identified as an important practice to minimise nutrient losses to the environment and maximise grass production. This is reported in binary (yes/no) format.
- **Protected urea fertiliser use** is associated with lower greenhouse gas emissions compared to use of Calcium Ammonium Nitrate (CAN). Protected urea is also associated with lower ammonia emissions compared to conventional straight urea fertiliser formulations and allows for greater nitrogen recovery for agronomic purposes. The indicator reported is the proportion of chemical N applied in protected urea form.
- **Low emission slurry spreading** or LESS (trailing shoe, trailing hose or injection methods) increases nitrogen retained in slurry and reduces the need for chemical fertiliser, as well as reducing nitrogen losses to the environment. The indicator reported is the proportion of farm slurry applied using LESS techniques.
- **Milk recording** (the practice of keeping detailed records of individual cow performance) was identified as a key aspect of dairy farm management practice from which farms could build on and improve herd health performance, breeding and milk yield. This reported is in binary (yes/no) format.

Cattle and sheep innovation indicators

For sheep and drystock cattle systems a common set of innovation indicators was used. These are:

- **Discussion group membership** was selected as indicating the degree of interaction with extension services and farming peers. This is reported in binary (yes/no) format.
- **Liming and Reseeding** were identified as important practices in grassland management. These indicators are reported in binary (yes/no) format.
- **Spring slurry spreading** (spreading at least 50% of total slurry between January and April) was identified as an important practice to minimise losses to the environment and maximise grass production. This is reported in binary (yes/no) format.
- **Protected urea fertiliser use** is associated with lower greenhouse gas emissions compared to Calcium Ammonium Nitrate (CAN). It is also associated with lower ammonia emissions compared to straight urea fertiliser formulations and greater nitrogen recovery for agronomic purposes. The reported indicator is the proportion of chemical N applied in protected urea form.
- **Low emission slurry spreading** or LESS (trailing shoe, trailing hose or injection methods) increases nitrogen retained in slurry and reduce the need for chemical fertiliser, as well as reducing nitrogen losses to the environment. The indicator reported is the proportion of farm slurry applied using LESS techniques.

Tillage innovation indicators

- ***Discussion group membership*** was selected as indicating the degree of interaction with extension services and farming peers. This is reported in binary (yes/no) format.
- ***Liming*** was identified as important practices in arable production. This is reported in binary (yes/no) format.
- ***Growing a main break crop*** (oilseed rape, peas, beans, linseed) was identified as best practice for tillage farms for disease and pest control. This is reported in binary (yes/no) format.

4 Interpretation of Sustainability Indicator Results

The main diagrams used to represent sustainability indicator results are provided below. Boxplots are used to display continuous data and allow the visualisation of the statistical distribution of the results for the population represented. The boxplots used here show the **10th, 25th, 50th, 75th and 90th percentiles** of the NFS sample’s population weighted distribution. An annotated **hypothetical example** is shown in Figure 4-1 below, using data on gross margin per hectare for dairy farms. The value of the percentiles reflect the distribution of results. For example, the 50th percentile (the median) in Figure 4-1 lies at approximately €1,400 per hectare, meaning that 50% of farms had a gross margin per hectare below this value (and conversely, 50% of farms had a gross margin per hectare greater than this value). A shorter range between percentiles indicates farms within this range have similar levels of performance. In the hypothetical dairy example below, the distance between the 90th and 75th percentiles is greater than the distance between the 50th and 75th percentiles, indicating that a larger number of dairy farms were closer to this central range, with a wider spread among farms earning significantly more.

For **indicators with binary scores**, bar charts show the proportion of farms that scored positively for the given indicator, as shown for dairy farm economic viability in Figure 4-2 below. To reflect how a given (non-economic) indicator relates to the economic performance of a farm, for most indicators, **farms are segmented** by performance into a **top, middle and bottom** performing third, on the basis of gross margin per hectare. This is also demonstrated in the example of Figure 4-2, where it can be seen in this hypothetical case that 88% of the top third of dairy farms, ranked by gross margin (GM) per hectare, were economically viable, compared to 34% for the bottom third.

Figure 4-1: Example Boxplot Gross Margin € per hectare





Figure 4-2: Example Bar Chart Proportion of farms



Dairy Farm Sustainability 2022

Key Messages



Economic

Family Farm Income reached €2,300 per ha



Environmental

GHG emissions per ha fell by 0.2 tonnes



Social

Just 4% of households were considered vulnerable




5 Dairy Farm Sustainability 2022

Economic Sustainability Indicators


In 2022, the average dairy farm **output per hectare** was €6,005, and the average market based **gross margin per hectare** was €3,509. Median values were slightly lower than the average, as shown in Figure 5-1.

Figure 5-1: Economic Return and Profitability of Land: Dairy Farms



Overall 93% of dairy farms were **economically viable** in 2022. This ranged from 99% for the top one third of economic performing dairy farms, to 85% for the bottom third, as illustrated in Figure 5-2.


Figure 5-2: Economic Viability: Dairy Farms



Average **income per labour unit** (unpaid family labour) for dairy farms in 2022 was €109,003. Average incomes per labour unit were €154,160, €110,424 and €61,621 for the top, middle and bottom performing farm


cohorts respectively. However, there was a large range in the return to family labour for dairy farms as shown in Figure 5-3.

Figure 5-3: Productivity of Labour: Dairy Farms




On average, dairy farms derived 94% of gross output directly from the market in 2022. The degree of **market orientation** was highest for the top third of dairy farms and the range was largest among the bottom third, as illustrated in Figure 5-4.

Figure 5-4: Market Orientation: Dairy Farms



For the full dairy farm population, there was a large range in family farm income per hectare across all three groups, as illustrated in Figure 5-5. The average family farm income per hectare on dairy farms was €2,332 in 2022. Within the farm profitability subcategories, the average income ranged from €3,364 per ha for the top performing cohort to €1,371 per ha for the bottom performers in economic terms.


Figure 5-5: Family Farm Income per hectare: Dairy Farms



Environmental Sustainability Indicators

Figure 5-6 indicates that the average dairy farm produced 606 tonnes of agricultural **GHG emissions** (in CO₂ equivalent) in 2022. It should be noted that this measure is based on the IPCC definition of agricultural emissions. At 73%, most dairy farm agricultural GHG emissions were associated with the production of milk output. A further 26.6% of dairy farm GHG emissions were allocated to beef production on these farms (this would include emissions from cull cows and calf sales and transfers). The remaining emissions, less than 1%, were associated with sheep production on dairy farms.


Figure 5-6: Agricultural GHG Emissions for the average Dairy Farm



The average dairy farm produced 9.44 tonnes of CO₂ equivalent GHG **emissions**


per hectare of UAA. The better performing dairy farms in an economic sense tended to operate at higher intensities and this is reflected in their higher emissions of GHG per hectare, as shown in Figure 5-7.

Figure 5-7: Agricultural GHG Emissions per hectare: Dairy Farms



When GHG emissions allocated to dairy production are expressed per kilogramme (kg) of milk output, the average dairy farm had GHG **emissions** of 0.84 kg CO₂ equivalent **per kg of milk produced.**² Figure 5-8 shows that those farms with a better economic performance also tended to have the lowest emissions intensity per kg of milk produced.

Figure 5-8: Agricultural GHG Emissions per kg of Milk: Dairy Farms




Emissions allocated to dairy output can also be expressed **per kg of fat and protein corrected milk (FPCM)**, which is standardized to 4% fat and 3.3% true protein

² Convert kg to litre by multiplying by 1.03


per kg of milk. The average farm had GHG emissions of 0.82 kg CO₂ equivalent per kg of FPCM produced in 2022. Figure 5-9 also shows that those farms with better economic performance also have lower emissions intensity per kg of FPCM produced.

Figure 5-9: Agricultural GHG Emissions per kg of FPCM: Dairy Farms



In 2022, the average dairy farm had **energy based GHG emissions** of 0.33 tonnes of CO₂ equivalent **per hectare**. In economic terms, the better performing dairy farms tended to operate at higher intensities, reflected in their higher emissions of energy based GHG per hectare, as shown in Figure 5-10.


Figure 5-10: Energy use related GHG Emissions per hectare: Dairy Farms



The average dairy farm's **energy based GHG emissions** were 0.0353 kg CO₂ equivalent **per kg of milk** in 2022. Figure 5-11 indicates that, similar to agricultural based GHG emissions intensity of milk production, lower energy based GHG emissions per kg of milk


produced is evident among farms with better economic performance.

Figure 5-11: Energy use related GHG Emissions per kg of Milk: Dairy Farms



The average dairy farm's energy based GHG emissions were 0.0331 kg CO₂ equivalent **per kg of FPCM** produced as shown in Figure 5-12. This indicator again shows that the top economic performers were more efficient in terms of FPCM produced per kg of energy related CO₂ emissions generated.

Figure 5-12: Energy GHG Emissions per kg of FPCM: Dairy Farms



Using the **LCA approach** (including both agricultural and energy based GHG emissions) the average dairy farm **carbon footprint of milk** was 1.06 kg CO₂ equivalent per kg of FPCM in 2022. Figure 5-13 again shows that lower emissions per kg of FPCM (on a LCA basis) was more prevalent among the group of top (economically) performing dairy farms.

Figure 5-13: Total LCA based GHG emissions (Agriculture & Energy) per kg of FPCM: Dairy Farms






Figure 5-14 indicates that the average dairy farm produced approximately 3 tonnes of ammonia (NH₃) emissions in 2022. This calculation is based on an approach consistent with the EPA national ammonia inventory methodology. The majority of dairy emissions (74%) were from milk based output, with 26% allocated to non-milk producing animal activities and a minor amount allocated to arable production.

Figure 5-14: Total Ammonia Emissions for the average Dairy Farm




The average dairy farm emitted 46.9 kg of NH₃ per hectare across the entire farm. Economically better performing farms tend to operate at higher intensities and this is reflected in higher emission of ammonia per hectare, as shown in Figure 5-15.

Figure 5-15: Ammonia Emissions kg per hectare: Dairy Farms




The average dairy farm emitted 0.004 kg of NH₃ per kg of FPCM produced. Figure 5-16 again shows that the top economic performing dairy farms produced milk at a lower NH₃ emissions intensity compared to the middle and bottom cohorts.

Figure 5-16: Ammonia Emissions per kg of FPCM: Dairy Farms



This result was replicated in the outcome on a kg of milk output basis, as shown in Figure 5-17. However, the NH₃ emissions per kg of milk was slightly higher at 0.0043.


Figure 5-17: Ammonia Emissions per kg of Milk: Dairy Farms



Nitrogen balance (excess of N inputs over outputs) averaged 158.6 kg N surplus per hectare across all dairy farms in 2022.


Figure 5-18 indicates that higher N surpluses per hectare are associated with superior economic performance. This is due to the greater production intensity on economically better performing farms.

Figure 5-18: N Balance per ha: Dairy Farms




The average dairy farm had an **N use efficiency** (NUE) of 27.7% in 2022. Figure 5-15 demonstrates that a slightly higher NUE was evident among the better economic performing farmers, with the largest range prevalent among the bottom cohort.

Figure 5-19: N Use Efficiency: Dairy Farms




Phosphorus balance (excess of inputs over outputs) averaged 10.4 kg P surplus per hectare across all dairy farms in 2022. Figure 5-20 shows that there was a larger range of results, especially for the top and middle performing cohorts.

Figure 5-20: P Balance per ha: Dairy Farms



The average dairy farm had a **P use efficiency** of 60.8%. Figure 5-21 indicates higher P use efficiency was more prevalent among the better economic performing farms.


Figure 5-21: P Use Efficiency: Dairy Farms



Social Sustainability Indicators


A minority of all dairy farm households, 4%, fell into the **vulnerable household** category (economically non-viable farm business and no off-farm employment). Figure 5-22 shows that there was a considerably larger proportion of households at risk among those farms with the lowest gross margin per hectare (10% among bottom third).

Figure 5-22: Household Vulnerability: Dairy




Overall, 83% of dairy farmers had received some formal agricultural education. Figure 5-23 shows that agricultural training rates were slightly higher across the middle and top performing cohorts.

Figure 5-23: Agricultural Education: Dairy




Only 6% of dairy **farmers live alone** and were thus classified as being at risk of isolation. Figure 5-24 indicates that the risk was lowest for the top economic performing cohort.

Figure 5-24: Isolation Risk: Dairy Farms




Across all dairy farms, 18% were identified as having a **high age profile**. shows that the percentage was slightly higher for the weaker economic performing dairy farms.

Figure 5-25: High Age Profile: Dairy Farms




On average, dairy farmers worked 2,498 hours per year on-farm (approximately 48 hours per week). Figure 5-26 shows that the number of hours worked was highest for the top and middle performing cohorts by economic performance. However, this figure does not take into consideration off-farm employment, or the share of hours worked by hired staff or other family members.

Figure 5-26: Hours Worked on farm: Dairy Farm Operator



On average, dairy farmers worked 2,657 hours per year between on and off-farm work (approximately 50.7 hours per week). Figure 5-27 shows that hours worked was slightly higher for the middle and bottom performing cohorts by economic performance.

Figure 5-27: Total Hours Worked: Dairy Farm Operator




Dairy Innovation Indicators

The innovation indicators analysed for dairy farms were, the use of **milk recording**, **membership** of a **dairy discussion group**, whether at least 50% of **slurry application** occurred the period January to April, use of **low emission slurry spreading** equipment, application of **protected urea fertiliser**, as well as **liming** & **grassland reseeding** rates.

Figure 5-28 shows that those farms with better economic performance were more likely to use **milk recording**. Over 62% of the dairy farmers in the top group were milk recording, compared to 41% in the bottom group.


Figure 5-28: Milk Recording: Dairy Farms



Better economic performance was more prevalent among **discussion group members**. Membership rates were higher

across the top economic performing group, at 55%, compared to 29% in the bottom cohort, as shown in Figure 5-29.

Figure 5-29: Discussion Group: Dairy Farms



The **application** of the majority of **slurry** in early spring was slightly higher across the top performing cohort at 63%, as shown in Figure 5-30. The middle and bottom cohorts had slightly lower level of spring time slurry application at 49% and 51% respectively.

Figure 5-30: Spring Slurry: Dairy Farms





Figure 5-31 illustrates the volume of slurry applied by **low emissions slurry spreading** equipment. On the average dairy farm, nearly 75% of all slurry applied by dairy farmers was via low emission slurry spreading methods. This ranged from 93% for the top performing cohort to 57% for the bottom performing cohort.

Figure 5-31: Slurry applied by Low emissions slurry spreading methods: Dairy Farms



The percentage of total chemical nitrogen applied in the form of **protected urea** averaged 14% across all dairy farms. This ranged from 18% for the top performing cohorts to 9% for the bottom group as illustrated by Figure 5-32.

Figure 5-32 Protected Urea Use: Dairy Farms




Figure 5-33 shows that **liming** was more prevalent among the better economic performers, with 58% of the top performing group engaging in this practice in 2022, compared to 46% for the bottom group.

Figure 5-33: Liming: Dairy Farms





Figure 5-34 shows that **reseeding** was also more common among the better economic performing farms. A higher percentage of farmers in the top and middle groups (34-35%) engaged in reseeding of grassland compared to the bottom group (22%) in 2022.

Figure 5-34: Reseeding: Dairy Farms



Cattle Farm Sustainability 2022

Key Messages



Economic

Family Farm Income was €370 per ha



Environmental

GHG emissions per ha fell by 0.3 tonnes



Social

About 1/3 of households were considered vulnerable




6 Cattle Farm Sustainability 2022

Cattle farms include both cattle rearing (mainly suckler based) and cattle finishing systems. Results for sustainability indicators for these systems in 2022 are presented below.

Economic Sustainability Indicators


The average **output per hectare** for cattle farms was €1,740, and the average gross margin per hectare was €769 in 2022. There was a large range in farm economic performance, as shown in Figure 6-1.

Figure 6-1: Economic Return and Profitability of Land: Cattle Farms




Only 27% of all cattle farms in the Teagasc NFS were defined as **economically viable**. As illustrated in Figure 6-2, the proportions deemed viable were 54%, 23% and 6% for the top, middle and bottom cohorts of farms by economic performance respectively.

Figure 6-2: Economic Viability: Cattle Farms




Across all cattle farms, the average **income per labour unit** was €17,154 in 2022. Figure 6-3 shows that this distribution was skewed by the top third of farms, which included a large number of relatively higher earners, with a mean income per labour unit of €32,681, compared with €15,345 and €3,452 for the middle and bottom cohorts of cattle farms respectively.

Figure 6-3: Productivity of Labour: Cattle




Market based output accounted for 70% of gross output across all cattle farms, with the remaining 30% accounted for by direct payment receipts. Figure 6-4 shows greater market orientation was exhibited across farms with better economic performance.

Figure 6-4: Market Orientation: Cattle Farms



The average family farm **income per hectare** on cattle farms was €372 in 2022. Across the subgroups, the average ranged from €718 for the top performing cohort to €71 for the bottom performers economically. Figure 6-5 shows significant ranges in income per hectare within and across the three groups, with a negative income per hectare returned by a section of the bottom performing cohort.


Figure 6-5: Family Farm Income per hectare: Cattle Farms



Environmental Sustainability Indicators


The average **cattle farm** produced 150.9 tonnes CO₂ equivalent of agricultural **GHG emissions** in 2022. Figure 6-6 shows that beef production was the principal source, generating 97% of these emissions. Sheep production was responsible for approximately 2.6% of total emissions on Irish cattle farms, and a very small proportion (less than 0.2%) was derived from other enterprises on these farms.

Figure 6-6: Agricultural GHG Emissions for the average Cattle Farm



The average cattle farm emitted 4.41 tonnes of CO₂ equivalent of agriculturally generated **GHG emissions per hectare** in 2022. Emissions per hectare were higher for the more profitable cattle farms, which tended to be stocked at a higher intensity.


Figure 6-7: Agricultural GHG Emissions per hectare: Cattle Farms



The **emissions** generated by cattle can be expressed in terms of their **live-weight output** (estimated using CSO price data). Figure 6-8 illustrates that there is a large range of **emissions per kg of beef live-weight** output. A positive association exists between emissions efficiency and economic performance. The top performing third of farms emitted, on average, 7.9 kg CO₂ equivalent per kg of live-weight beef produced, compared with 11.2 kg for the bottom performing third of cattle farms. The average level of GHG emissions across all


farms was 9.4 kg CO₂ equivalent per kg beef of live-weight produced.

Figure 6-8: Agricultural GHG Emissions per kg live-weight beef produced: Cattle Farms




The average cattle farm emitted 0.18 tonnes of CO₂ equivalent of **energy based GHG emissions per hectare** in 2022, as illustrated in Figure 6-9. Emissions per hectare were higher for the more profitable cattle farms, which tended to be stocked at a higher intensity.

Figure 6-9: Energy GHG Emissions per hectare: Cattle Farms




On average, **energy based GHG emissions** across all cattle farms was 0.39 kg of CO₂ equivalent **per kg beef live-weight** produced. Figure 6-10 illustrates that energy based GHG emissions per unit of product were also lower on farms with better economic performance. The top third produced an average of 0.33 kg CO₂ energy-based emissions per kg of live-weight beef produced, while for the bottom performing third this figure was 0.48 kg.

Figure 6-10: Energy use related GHG Emissions per kg live-weight beef: Cattle Farms



The average **cattle farm** emitted 0.76 tonnes of **ammonia (NH₃)** in 2022. Over 98% of total NH₃ emissions were linked to beef production, the remainder reflected emissions from a sheep and tillage enterprise on cattle farms, as shown by Figure 6-11.

Figure 6-11: Total Ammonia Emissions for the average Cattle Farm



On average, cattle farms emitted 22 kg of **NH₃ per hectare** in 2022. This ranged from 28.3 kg per hectare for the top performing cohort, to 17 kg per hectare for the bottom third, as shown by Figure 6-12. Emissions per hectare were higher for the more profitable cattle farms, which also tend to be stocked at a higher intensity.

Figure 6-12: Ammonia Emissions per hectare: Cattle Farms




Figure 6-13 illustrates that, in terms of **live-weight** of beef produced, the more profitable cattle farmers have a lower level of **ammonia emissions**. There was a large range within the results, especially for the bottom performing cohort of cattle farmers. On average, a kg of live-weight beef was produced at an intensity of 0.0499 kg of NH₃.

Figure 6-13: Ammonia Emissions per kg live-weight beef produced: Cattle Farms






Figure 6-14 indicates that the **nitrogen surplus per hectare** tended to be higher on cattle farms that performed better economically. In general, these farms are operated more intensively. The top performing third of cattle farms had an average nitrogen surplus of 59.3 kg N per hectare, compared to 31.6 kg N per hectare for the bottom third of farms.

Figure 6-14: N Balance per ha: Cattle Farms




The average **N use efficiency (NUE)** across all cattle farms was 33.9%, but the range in NUE across the sample of cattle farms was significant, as shown in Figure 6-15. Despite the higher application rates, NUE tended to be higher across the middle and top economic performing cohorts.

Figure 6-15: N Use Efficiency: Cattle Farms




At the farm gate boundary, the **P surplus** across all cattle farms averaged 2.4 kg of P per hectare. There was a large range in P surpluses, especially across the better performing farms economically, as shown in Figure 6-16.

Figure 6-16: P Balance per ha: Cattle Farms



At the farm gate boundary, the average farm **P use efficiency** (PUE) across all cattle farms was 89.2%. Figure 6-17 shows that higher PUE was again more prevalent on farms that performed better in economic terms. Average PUE ranged from 90.6% for the top third to 72.8% for the bottom third of cattle farmers.

Figure 6-17: P Use Efficiency: Cattle Farms



Social Sustainability Indicators

Overall, 33% of all cattle farm households were considered **vulnerable** (a non-viable farm business with no off-farm employment). Figure 6-18 confirms that this vulnerability was associated with weaker economic performance, with 29% and 50% of the middle and bottom third of farms deemed vulnerable, compared to 18% of the top third.

Figure 6-18: Household Vulnerability: Cattle






Figure 6-19 indicates that **educational attainment** was positively associated with the better economic performing farms. A total of 49% of cattle farmers had some level of formal agricultural education.

Figure 6-19: Agricultural Education: Cattle Farms




Overall, 17% of cattle farm operators were classified as being at **risk of isolation**; i.e. where the farmer lives alone. This was especially prevalent among farms in the lower profitability cohorts, where 23% of farmers live alone, as shown in Figure 6-20.

Figure 6-20: Isolation Risk: Cattle Farms



Additionally, 41% of cattle farms were classified as having a **high age profile**. High age profile was lowest for the middle cohort as shown in Figure 6-21.

Figure 6-21: High Age Profile: Cattle Farms



The average cattle farm operator had 1,370 **hours worked on farm** over the year (an average of 26.3 hours per week). The top economically performing cohort worked on average of 1,507 hours on farm compared to 1,255 and 1,345 for middle and bottom groups as outlined in Figure 6-23.

Figure 6-22: Hours Worked on Farm: Cattle Farm Operator





Figure 6-23 shows total **hours worked on and off-farm** was slightly higher for the middle and top compared to the bottom cohort. On average, cattle farmers worked 2,139 hours in 2022 between on and off-farm work (approximately 41 hours per week).

Figure 6-23: Total Hours Worked: Cattle Farm Operator




Cattle Farm Innovation Indicators

Six innovation indicators were examined for cattle farms: whether at least 50% of **slurry application** occurred in the period January to April, the proportion of slurry applied using **low emission slurry spreading** equipment, proportion of chemical applied in the form of **protected urea** fertiliser, application of **lime**, grassland **reseeding** and whether the farmer was a member of a **discussion group**.

Figure 6-24 shows that those in the top and middle economic performing group applied a lot more **slurry application** in springtime


(46-53%) compared to the bottom (32%) cohort.

Figure 6-24: Spring Slurry Application: Cattle Farms



On average, nearly 34% of all slurry applied by cattle farmers was via **low emission slurry spreading** methods. This ranged from 38-40% for the top & middle performing cohort to 24% for the bottom performing cohort, as shown in Figure 6-25.

Figure 6-25: Low emission slurry spreading: Cattle Farms



The percentage of total chemical nitrogen applied in the form of **protected urea** averaged 4.3% across all cattle farms in 2022. This ranged from 9.4% for the top performing cohorts to 1% for the bottom group as illustrated in Figure 6-26.

Figure 6-26: Protected Urea use: Cattle Farms




Figure 6-27 shows that **liming** rates were higher for the top performing cattle farm, at 35%, compared to 16% for the bottom cohort.

Figure 6-27: Liming: Cattle Farms






Figure 6-28 shows that 23% of the top economic performing cohort were members of a **discussion group**, compared to 12% and 6% in the middle and bottom cohort respectively.

Figure 6-28: Discussion Group: Cattle Farms



Reseeding levels ranged from 16% for the top cohorts to 8-9% for the middle and bottom performing cohort as shown in Figure 6-29.

Figure 6-29: Re-seeding: Cattle Farms



Sheep Farm Sustainability 2022

Key Messages



Economic

Family Farm Income fell to €340 per ha



Environmental

GHG emission fell by 0.8 tonnes per ha



Social

Just under 1/3 of households were considered vulnerable




7 Sheep Farm Sustainability 2022

Economic Sustainability Indicators


The average **gross output per hectare** for sheep farms was €1,475 in 2022, and the average **gross margin** was €625 per hectare.

Figure 7-1: Economic Return and Profitability of Land: Sheep Farms



Across all sheep farms, 26% were defined as economically **viable**. Figure 7-2 shows that, ranked by economic performance, the proportion of viable sheep farms ranged from 48% for the top third to 12% for the bottom third of farms.


Figure 7-2: Economic Viability: Sheep Farms



The average **income per unpaid labour unit** on sheep farms was €14,890. In common with cattle farms, there was a large range in economic performance, with the top third of sheep farms earning a mean income per labour unit of €25,827, compared with


€7,204 for the bottom third (see Figure 7-3). Median income for the three cohorts was €20,111, €10,688 and €3,969 respectively.

Figure 7-3: Productivity of Labour: Sheep Farms




For the average sheep farm, approximately 65% of output was generated from the market, with the remaining 36% derived from direct payments. Figure 7-4 indicates that **market orientation** was positively associated with economic performance, with the top third of farms, based on economic performance, producing 76% of output from the market, compared with just over 49% on average for bottom third. Figure 7-4 also indicates a significant range in market orientation across the bottom performing cohort in particular.

Figure 7-4: Market Orientation: Sheep Farms



The average **family farm income per hectare** on sheep farms was €340 in 2022. Across the subgroups, this average ranged from €641 for the top performing cohort to €68 for the bottom performers economically. Figure 7-5 shows significant ranges in income per hectare across the three groups.


Figure 7-5: Family Farm Income per hectare: Sheep Farms



Environmental Sustainability Indicators


In 2022, the average **sheep farm** produced 136.2 tonnes CO₂ equivalent of agricultural **GHG emissions**. Figure 7-6 indicates that 54.4% of these emissions were generated by the sheep enterprise, with the remaining emissions (45%) generated by a cattle enterprise present on specialist sheep farms, with the remainder coming from other sources (minor arable enterprise).

Figure 7-6: Agricultural GHG Emissions for the average Sheep Farms




On average, sheep farms produced **GHG emissions** of 3.3 tonnes of CO₂ equivalent **per hectare**. Higher emissions per hectare were associated with the more profitable sheep farms, as shown in Figure 7-7. However, there was a large range of results.

Figure 7-7: Agricultural GHG Emissions per hectare: Sheep Farms




The **GHG emissions** generated by sheep are shown **per kg of live-weight** output produced (estimated using CSO price data). Figure 7-8 shows that the emissions intensity per kg of live-weight produced were negatively associated with economic performance. The top and middle third of farms generated 6.9 to 9.4 kg CO₂ equivalent per kg live weight produced respectively, compared to 11.6 kg CO₂ equivalent for the bottom cohorts on average. There was a noticeably large range in the emissions per kg of live-weight across the farms in the bottom cohort.

Figure 7-8: Agricultural GHG Emissions per kg live-weight produced: Sheep Farms




The average sheep farms had **energy related GHG emissions** of 0.16 tonnes CO₂ equivalent **per hectare**. Higher emissions per hectare were associated with the more profitable sheep farms, as shown in Figure 7-9.

Figure 7-9: Energy GHG Emissions per hectare: Sheep Farms




Better economic performance was also linked with lower **energy related GHG emissions per unit of output**, as shown in Figure 7-10. The bottom third of farms in economic terms emitted 0.80 kg CO₂ equivalent per kg live-weight sheep meat produced from energy based emissions, compared to 0.37 and 0.45 kg CO₂ for the top and middle third respectively.

Figure 7-10: Energy use related GHG Emissions per kg live-weight produced: Sheep Farms




On average, specialist **sheep farms** had 0.56 tonnes of **NH₃ emissions** in 2022. Even though the main output on these farms is sheep based, the majority of the NH₃ emissions related to cattle production (64%), with 35% relating to sheep production. The remaining residual portion related to production of tillage crops.

Figure 7-11: Total Ammonia Emissions for the average Sheep Farm



On average, a specialist sheep farm emitted 12.7 kg of **NH₃ per hectare** in 2022. Higher per hectare emissions were associated with economically better performing farms as shown in Figure 7-12. These farms tend to operate at a higher stocking intensity.


Figure 7-12: Ammonia Emissions per hectare: Sheep Farms



Lower ammonia emissions intensity of production was again more common among the better economically performing sheep farms. Farms in the top and middle performing cohort in economic terms were found to produce a kg of live-weight sheep


meat with a lower NH₃ emission footprint, as shown in Figure 7-13. On average, sheep farmers produced 0.022 kg of NH₃ emissions per kg of live-weight sheep meat.

Figure 7-13: Ammonia Emissions per kg live-weight produced: Sheep Farms



As with cattle farms, the sheep farm based **nitrogen surplus per hectare** was positively associated with economic performance, due to greater production intensity on the more profitable sheep farms (as shown in Figure 7-14). The top third of farms, ranked by gross margin per hectare, had an average nitrogen surplus of 44.3 kg per hectare, compared with 30.6 and 15.6 kg per hectare for the middle and bottom cohorts respectively.


Figure 7-14: N Balance per ha: Sheep Farms



The average **N use efficiency** (NUE) across all sheep farms was 37.6%. Higher NUE was again associated with better economic performance, as shown in


Figure 7-15.

Figure 7-15: N Use Efficiency: Sheep Farms




P balances across all specialist sheep farms were 2 to 4 kg per ha on average. There was a large range of results across the three cohorts, especially the top performing group, as shown by Figure 7-16.

Figure 7-16: P Balance per ha: Sheep Farms



Farm gate level **P use efficiency** (PUE) averaged 78.3% across all sheep farms in 2022. Figure 7-17 shows that higher PUE was associated with farms with better economic performance.


Figure 7-17: P use efficiency: Sheep Farms



Social Sustainability Indicators


Over 29% of all sheep farm households were considered **vulnerable** in 2022. Figure 7-18 shows that this ranged from 10% for the top performing sheep farms to 39% for the bottom group.

Figure 7-18: Household Vulnerability: Sheep Farms



Overall, 60% of sheep farmers had received formal **agricultural education**. Figure 7-19 shows that agricultural education was higher among the top and middle third of farms when ranked by economic performance.

Figure 7-19: Agricultural Education: Sheep Farms



On average, 13% of all specialist sheep farms were classified as being at **risk of isolation**. Figure 7-20 shows that this was significantly higher among the bottom performing cohort of sheep farms at 20%.

Figure 7-20: Isolation Risk: Sheep Farms






Figure 7-21 shows that the proportion of all specialist sheep farms with a high age profile was 28%. The bottom performing group had the highest age profile on average.

Figure 7-21: High Age Profile: Sheep Farms




Sheep farmers had an average of 1,477 **hours worked on farm** per year in 2022 (or 28.4 hours a week). The top and bottom performing cohort tend to work the most hours on farm at 1,508 to 1,514 hours, compared to the middle group at 1,407 hours (Figure 7-22).

Figure 7-22: Hours Worked On Farm: Sheep Farm Operators



On average, in 2022, sheep farmers had 2,246 **hours worked on and off-farm work** (approximately 43.2 hours per week). Figure 7-23 shows that total hours worked was lower across the middle cohort by economic performance.

Figure 7-23: Total Hours Worked: Sheep Farm Operator




Sheep Farm Innovation Indicators

The five innovation indicators selected for sheep farms were whether at least 50% of **slurry application** occurred in the period January to April, the portion of chemical N fertiliser applied in the form of **protected urea**, application of **lime**, grassland **reseeding** and whether or not the farm operator was a member of a **discussion group**.

Figure 7-24 shows that those in the top and middle economic performing group (23 to 26%) undertook more **slurry application** in the springtime, compared to the bottom cohort (19%). However, it should be noted that sheep farms tend to be more associated with farmyard manure (i.e. solid) type storage systems, which do not lend themselves to early season application.

Figure 7-24: Spring Slurry: Sheep Farms



As illustrated in Figure 7-25, the use of **protected urea** fertiliser by sheep farmers was very limited. On average only 2% of chemical N fertiliser applied was in the form of protected urea in 2022.

Figure 7-25: Protected Urea use: Sheep Farms




Figure 7-26 shows that **liming** activity was again more prevalent across the better economic performing farms, with 35% of the middle performing cohort by economic performance engaged in liming, compared to 12% of the bottom group.

Figure 7-26: Liming: Sheep Farms




Figure 7-27 shows that higher levels of **reseeding** were associated with the sheep farms that performed better in economic terms.

Figure 7-27: Reseeding: Sheep Farms





Figure 7-28 shows that membership of a **discussion group** was higher (24%) among the top cohorts versus 10% for bottom group.

Figure 7-28: Discussion Group: Sheep Farms



Tillage Farm Sustainability 2022

Key Messages



Economic

Family Farm Income reached over €1,000 per ha



Environmental

GHG emissions fell by 0.3 tonnes per ha



Social

Just 10% of households were considered vulnerable




8 Tillage Farm Sustainability 2022

Economic Sustainability Indicators


The average **gross output** and **gross margin per hectare** for tillage farms was €2,812 and €1,581 respectively in 2022. However, there was a large distribution around the average, as illustrated by Figure 8-1.

Figure 8-1: Economic Return and Profitability of Land: Tillage Farms



In 2022, 79% of tillage farms were classified as economically **viable**. Figure 8-2 shows that the bottom group had lower levels of viability, at 63% compared to 90% for the top performing group.


Figure 8-2: Economic Viability: Tillage Farms



The average tillage farm **income per unpaid labour unit** (for unpaid family labour) was €97,910 in 2022. Figure 8-3 shows that there is a large range in incomes on tillage farms, with the top one-third (ranked by gross


margin per hectare) earning significantly more than the middle and bottom cohorts per labour unit provided. For some of the most profitable tillage farms, income per family labour unit is especially high, due to the large proportion of the labour utilised on tillage farms being supplied by hired labour (via the use of external contractors).

Figure 8-3: Productivity of Labour: Tillage Farms




In 2022, tillage farms generated 82% of their output value from the market on average. Figure 8-4 shows that the top third of tillage farms had a **market orientation** of 85% compared to 80% for the bottom group on average.

Figure 8-4: Market Orientation: Tillage Farms



The **average family farm income per hectare** on tillage farms was €1,043 in 2022. Median income ranged from €1,434 from the top performing cohort to €650 for the bottom performers economically. Figure 8-5 shows significant ranges in income per hectare across the three groups.


Figure 8-5: Family Farm Income per hectare: Tillage Farms



Environmental Sustainability Indicators


The average **tillage farm** produced 137.5 tonnes CO₂ equivalent of agricultural **GHG emissions** in 2022 as illustrated in Figure 8-6. However, only 28.4% of GHG emissions on these farms were generated from crop production. Despite being specialised in crop production, 62.3% of tillage farm emissions were from cattle present on these farms, with a further 9.4% from sheep.

Figure 8-6: Agricultural GHG Emissions for the average Tillage Farm




The average specialist tillage farm had agricultural **GHG emissions** of 1.9 tonnes CO₂ equivalent **per hectare** in 2022. Emissions per hectare tended to be lower for the better performing economic cohorts as illustrated by Figure 8-7.

Figure 8-7: Agricultural GHG Emissions per hectare: Tillage Farms




Specialist tillage farms on average produced 0.21 tonnes of **energy based GHG emissions per hectare** in 2022. Higher emissions per hectare were associated with higher economic performance as illustrated in Figure 8-8.

Figure 8-8: Energy GHG Emissions per hectare: Tillage Farms




Tillage farms on average had **NH₃ emissions** of 0.72 tonnes in 2022. Again, even though the main farm output on such farms is crop related, the bulk of NH₃ emissions are associated with cattle rearing, at 59%. Of the remaining emissions, 38% were associated with tillage production and 3% with a sheep enterprise.

Figure 8-9: Total Ammonia Emissions for the average Tillage Farm




The average specialist tillage farm emitted 9.1 kg of **NH₃ per hectare** in 2022. Emissions per hectare were highest for the top and bottom cohorts.

Figure 8-10: Total Ammonia Emissions per hectare: Tillage Farms




The average **N surplus** was 40.5 kg per hectare, but there was a large range in the farm results as seen in Figure 8-11. The bottom cohort indicated higher N surpluses per hectare on average.

Figure 8-11: N Balance per hectare: Tillage Farms




Across all tillage farms, the average **N use efficiency (NUE)** was 66.6%. There was a large distribution in NUE across the three groups as illustrated in Figure 8-12.

Figure 8-12: N Use Efficiency: Tillage Farms




The average **P balance** across all tillage farms was 3.6 kg per hectare. However, as illustrated in Figure 8-13, there was again a large range of results around these group averages. Better farms, in economic terms, tended to have slightly lower P balances.

Figure 8-13: P Balance per hectare: Tillage Farms



P use efficiency (PUE) averaged 90.4% across all tillage farms. PUE tended to be higher across the top performing group, compared to the middle and bottom cohorts, as illustrated by Figure 8-14.


Figure 8-14: P Use Efficiency: Tillage Farms



Social Sustainability Indicators


On average, 10% of tillage farm households are considered economically **vulnerable**. Figure 8-15 indicates that household vulnerability was highest across the bottom cohort, at 21%. None of the top performing cohort were classified as vulnerable.

Figure 8-15: Household Vulnerability: Tillage




A total of 70% of tillage farmers had received some level of formal **agricultural education** or training. Figure 8-16 shows that this rate was higher for the better performing tillage farms economically.

Figure 8-16: Agricultural Education: Tillage Farms




Overall, 18% of tillage farms were identified as being at **risk of isolation** (i.e. where the farm operator lived alone). At 28%, this rate was highest for the bottom performing cohort, as illustrated by Figure 8-17.

Figure 8-17: Isolation Risk: Tillage Farms




An average of 25% of tillage farms were identified as having a **high age profile**. Figure 8-18 shows that over 41% of farm households in the bottom group had a high age profile.

Figure 8-18: High Age Profile: Tillage Farms




The average tillage farmer had 1,416 **hours worked on farm** in 2022 (27.2 hours per week). However, Figure 8-19 shows that there is significant variables in results across the different cohorts.

Figure 8-19: Hours Worked on Farm: Tillage Farms



On average, tillage farmers had 2,133 **hours worked on and off-farm** per year in 2022 (approximately 41 hours per week). Figure 8-20 shows that total hours worked tended to be higher across the middle cohort by economic performance.

Figure 8-20: Total Hours Worked: Tillage Farm Operator




Tillage Innovation Indicators

The innovation indicators examined for tillage farms were: **liming** rates, membership of a **discussion group** and growing of a **break crop**.

Figure 8-21 shows that **liming** rates were higher for the middle and top performing cohorts (27-33%) compared to the bottom (18%) performing cohort.

Figure 8-21: Liming: Tillage Farms



On average between 31% of tillage farms were in **discussion groups**. This ranged from 31-37% for the middle and top group compared to 24% for bottom performing cohort. However, this includes all types of discussion groups (e.g. beef and sheep).

Figure 8-23: Break Crops: Tillage




Figure 8-22: Discussion Group: Tillage Farms




Figure 8-23 shows that 27% of the top performing cohorts grew a **break crop** compared to 7-10% for the bottom and middle groups.

Farm System Comparisons 2022

Key Messages



Economic

Profitability on Dairy and Tillage farms increased, but remained unchanged on Beef and Sheep farms



Environmental

All farm systems showed a decline in GHG emissions per ha



Social

The rate of household vulnerability remained high on Beef and Sheep farms



9 Farm System Comparisons 2022

Economic Indicators

A comparison of economic sustainability indicators across different farm types is shown in Figure 9-1. Specialist dairy farms show the strongest economic performance, significantly ahead of all other systems in terms of gross output, gross margin and family farm income on a per hectare basis.


Output, Margins and Income: Tillage farms were ahead of both cattle and sheep farms (whose performance were relatively similar) in terms of gross output, gross margin and family farm income per hectare, but tillage farms were similar to dairy farms in terms of income per labour unit. Sheep and cattle farms returned significantly lower income per labour unit in comparison to dairy farms and tillage farms in 2022.

Market Orientation: The various farm systems are most similar in terms of market orientation, with dairy and tillage having the greatest share of gross output from the market.

Viability: Cattle and sheep farms are most at risk financially, with only 26% of sheep farms and 27% of cattle farms classed as economically viable in 2022. Dairy farms were the most economically viable (93%), followed by tillage systems (79%).

Caveat: It is important to note that these are average values for each farm type and that earlier analysis in this report has highlighted the range around these average values in the case of each farm system type. **Averages, while useful, do not tell the full story.** In some cases, the extent of the distribution around the average is such that there may be an overlap in the distribution of performance between different farm systems.

Figure 9-1: Economic Sustainability: Farm System Comparison 2022 (average per system)



Environmental Indicators

The environmental sustainability of farms is more difficult to compare directly across different farm systems, as the indicators are more directly linked with the type of farming undertaken and the different outputs produced. More detail can be obtained by comparing within farm type variations (see previous section), but some shared environmental indicators across different farm types are presented in Figure 9-2.

Greenhouse Gas Emissions: Animal based farming systems typically have higher greenhouse gas emissions per hectare than tillage systems, but this is to be expected due to the greater emissions associated with animal as opposed to crops production, especially in ruminant systems. Dairy farms show the highest emissions on a per hectare basis, significantly greater than any other system. This is attributable to the greater production intensity on these farms. Dairy based emissions per hectare are a function of higher stocking rates, more energy intensive diets for dairy cows and greater use of chemical fertilisers than is found in other livestock systems. In terms of kg of GHG emissions per euro of output generated, livestock farms (especially cattle) had much higher emissions due to the lower value of output generated in beef and sheep compared to dairy systems.


Ammonia Emissions: In common with GHG emissions, ammonia emissions per hectare were significantly higher on dairy farms compared to all

other systems. Cattle farms had the next highest level of emissions per hectare (though on average these were less than half those of the average dairy farms) followed by sheep and tillage farms. In terms of ammonia (NH₃) emissions per euro of market output generated, cattle farms exhibited the highest ammonia emissions intensity (due to the generally lower levels of output) followed by sheep farms. Tillage farms have the lowest level of ammonia emission per euro of output generated due to the low number of livestock on these farms on average.

Nitrogen use: Dairy farms have the largest N surplus per hectare due to the higher levels of livestock production intensity per hectare in this system. In terms of the input-output accounting NUE metric, dairying is more similar to the other livestock systems (but still lower), while tillage farms have greater NUE on average. It should be noted that tillage systems by their nature will have higher NUE, as the nitrogen is not cycling through an animal (and subject to the various loss pathways).

Phosphorus use: Dairy farms had the highest farm gate level P balances, significantly higher than those for the cattle, sheep and tillage systems. However, this metric should be interpreted with caution, as reference to a soil test is required to establish optimal P balance on farms and such soil test data are unavailable for farms in the NFS. PUE was highest on tillage farms, which was higher than that observed across all of the livestock systems.

Figure 9-2: Environmental Sustainability: Farm System Comparison 2022 (average per system)



Social Indicators

Comparison of the social sustainability indicators of different farm types (in

Figure 9-3) shows a similar overall trend to the economic performance indicators shown in Figure 9-1, with dairy and tillage farms being distinct from cattle and sheep systems, with respect to their social sustainability performance, but with some notable exceptions.


Hours Worked: The greater labour intensity of dairying is illustrated by the longer hours worked on farm. When accounting for total hours worked (on and off-farm employment), dairy farmers still have the highest number of hours worked on average, but the gap between dairy farms and other farm systems is reduced significantly.

Household Vulnerability: Given that there were lower levels of economic viability across cattle and sheep farms (see Figure 9-1) these systems were also more likely to have a more vulnerable household structure (non-viable farm business with no off-farm employment within the household).

High Age Profile: Cattle, sheep and then tillage farms were also more likely to have a high age profile, while cattle and tillage farms were more likely to be operated by farmers living alone. However, there was less variation within these measures than for other social sustainability indicators.

Agricultural Education: On average, dairy farmers were slightly more likely to have received agricultural education or training compared to other farm systems.

Figure 9-3: Social Sustainability: Farm System Comparison 2022 (average per system)



Time Series Comparisons by Farm System

Key Messages



Economic

The gap between dairy and tillage incomes relative to beef and sheep incomes is increasing



Environmental

A slowly improving trend in emerging in some environmental indicators



Social

Household vulnerability is falling across all farm systems, although the level remain high on Beef and Sheep farms

10 Time Series Comparisons with a three year rolling average: 2017-2022


Building on research presented in previously published Teagasc Sustainability reports (Hennessy et al., 2013; Lynch et al., 2016; Buckley et al., 2019; 2020a; 2020b; 2021), the evolution of farm-level sustainability indicators can be tracked over time. The figures presented below highlight changes in indicator scores, with averages presented across all farm types. As short term input and output price volatility and weather events in a given year can occur and distort intertemporal trends, results below are presented on the basis of a **three year rolling average** (i.e. the result for 2017 is based on the average of the years 2015 to 2017 inclusive and is labelled as such). For reference, the **annual average** results for each indicator are also provided in **Appendix 1**.

It is important to appreciate that some factors influencing the various indicator measures shown here are partially **within the control of an individual farmer** (e.g. input use efficiency) and hence may be improved by changes in farmer behaviour, while others factors are **outside of the control of an individual farmer** (e.g. farm output prices, weather conditions, soil quality). Since farming is influenced by **weather conditions**, which vary from year to year, and which therefore may affect the level of production or the level of input utilisation in a given year, this limits the inferences that can be drawn from one year movements in such time series. The reported annual data contain both the signal and noise components and the use of the three year moving average based indicators allows for the signal component of the indicator to be more apparent.

10.1 Economic sustainability indicators


Figure 10-1 shows that the value of economic return to land (gross output (€) per hectare) tended to increase over the study period. However, across individual farm systems, there are notable differences. Dairy farms have significantly higher levels of output per hectare compared to all other systems, especially towards the end of the study period. Tillage farmers were next highest, ahead of cattle and sheep systems. Additionally over the period studied the rate of growth in output value per hectare on dairy farms was considerably higher than on all other farm types. It should be noted that some of the increase observed, particularly on dairy farms relates to the increase in the price of milk rather than the volume of output produced.

Figure 10-1: Economic Returns to Land: 3 year rolling average 2017-2022



The profitability of land (gross margin per hectare) in dairying is again significantly higher than for all other systems and tended to increase over the years, significantly so at the end of the study period. Tillage farms again have the second highest gross margin per hectare. The lowest gross margins per hectare are on cattle and sheep farms, as illustrated in Figure 10-2.

Figure 10-2: Profitability of Land: 3 year rolling average 2017-2022



The ranking of systems based on Family Farm Income per hectare mirrors the ranking for profitability of land, with dairy incomes significantly higher than for all other systems. Similarly, tillage farms are again ranked second. The lowest family farm income per hectare are on cattle and sheep farms, as illustrated by Figure 10-3. Again family farm income on dairy farms was seen to significantly increase towards the end of the study period.

Figure 10-3: Family Farm income: 3 year rolling average 2017-2022




Figure 10-4 illustrates that farm income per unpaid labour unit broadly follows a similar trend to the gross output, gross margin and family farm income per hectare indicators. However, the differences between farm types when income per family labour unit is considered are less pronounced relative to the differences in gross output, gross margin and family farm income per hectare. This is due to the adjustment made to reflect different labour intensities of each production system. Returns to labour are significantly higher on dairy and tillage farms, compared to cattle and sheep systems.

Figure 10-4: Productivity of Labour: 3 year rolling average 2017-2022






Figure 10-5 illustrates that dairying is the most market orientated of all the systems (88 to 92%) followed by tillage systems (74 to 78%). The market orientation of cattle and sheep systems was the lowest at between 59% and 66%.

Figure 10-5: Percentage of Output Derived from Market: 3 year rolling average 2017-2022



The same trends over time are also observed in terms of farm economic viability. Dairy and tillage systems have significantly higher levels of viability compared to cattle or sheep farms over the period examined. Viability, as with the other economic indicators, is effected by variations in sectoral input and output prices over the period.

Figure 10-6: Economic Viability: 3 year rolling average 2017-2022



10.2 Environmental sustainability indicators

Figure 10-7 shows that the overall farm average agricultural GHG emissions per hectare have stabilised over the study period (5.1 tonnes CO₂ equivalent per hectare). Due to the more intensive nature of production in dairy systems compared to all other grassland systems, agricultural GHG emissions per hectare on dairy farms are 2 to 4 times higher compared to other farm systems. The main trends observed are an increase in dairy emissions per hectare and the relative stability or reduction in emission intensity per hectare across the other farm systems.

Figure 10-7: Ag. Greenhouse Gas Emissions per hectare: 3 year rolling average 2017-2022




Figure 10-8 shows that energy based GHG emission have generally remained stable or declined slightly over the study period. Energy based emissions were highest on dairy farms, since they are greater users of fuel and electricity.

Figure 10-8: Energy Greenhouse Gas Emissions per hectare: 3 year rolling average 2017-2022




Figure 10-9 illustrates that emissions per euro of output generated are significantly higher across livestock rearing farms. These results are reflective of the greater value of output produced in dairying and the lower emissions associated with tillage. The increase in dairy emissions per hectare, shown in Figure 10-7, is not reflected in a similar evolution in the emissions per euro output here as output prices have increased significantly towards the end of the study period as illustrated in Figure 10-9. It should be noted that agricultural output prices have increased by over 48% between 2015 and 2022 (CSO, 2023b).

Figure 10-9: Ag. GHG Emissions per Euro output: 3 year rolling average 2017-2022




Figure 10-10 illustrates energy related GHG emissions per euro of market based gross output. Results follow a similar pattern to that of agricultural based emissions, where energy emissions per euro of output are significantly higher across cattle/sheep farms compared to dairying, over the period presented. Across all farm systems, energy emissions per euro of output showed a declining trend over the study period. Again it should be noted that agricultural output prices have increased by over 48% between 2015 and 2022 (CSO, 2023b).

Figure 10-10: Energy related GHG Emissions per Euro output: 3 year rolling average 2017-2022




Figure 10-11 illustrates that on a three year rolling average basis across all farms, ammonia (NH₃) emissions per hectare increased over the early part of the study period before levelling off and then decreased towards the end of the period. Again, due to the more intensive nature of production, NH₃ emissions per hectare are significantly higher for dairy systems compared to all other grassland systems and especially tillage.

Figure 10-11: Kg of Ammonia Emissions per hectare: 3 year rolling average 2017-2022





Figure 10-12 illustrates NH₃ emissions per euro of market based gross output. Results indicate that emissions per euro of output are higher on cattle and sheep farms compared to all other systems over the study period. This is a function of the low levels of output on these farms. Dairy had the second lower levels of NH₃ emissions per euro of output generated (due to high output value). Tillage farms had the lowest emissions per euro of market based output. Again, it should be noted that agricultural output prices have increased by over 48% between 2015 and 2022 (CSO, 2023b).

Figure 10-12: Ammonia (NH₃) Emissions per Euro Output: 3 year rolling average 2017-2022



Across all farm systems, the N balance per hectare was similar at the end versus the start of the period examined. Again, due to the more intensive nature of production, N surpluses are significantly higher for dairy farms compared to all other systems. Due to the non-livestock orientated nature of production on tillage farms, N surpluses are, on average, lowest across these farms over the period examined. N surpluses are affected by a range of factors, some within and some (such as variability in the weather) outside the farmer's control. Higher N surplus years tended to be associated with adverse weather conditions.

Figure 10-13: Nitrogen Balance per ha: 3 year rolling average 2017-2022




Figure 10-14 illustrates that P balances have tended to increase over the study period. P surpluses are significantly higher on dairy farms compared to all other systems. It should also be noted that farm gate P balances must be interpreted with care, since establishing the optimal balance requires a soil test. Farmers are allowed to run significant farm gate surpluses, if soil P status is sub optimal (deficient). In 2022, Teagasc analysed a total 38,143 soil samples comprising of dairy, drystock and tillage farm enterprises (Teagasc, 2022). Results indicate that 50% of samples taken are P deficient (at either index 1 or 2 for phosphorus).

Figure 10-14: Phosphorus (P) Balance per ha: 3 year rolling average 2017-2022




Figure 10-15 illustrates that dairy and cattle farms tended to have the lowest NUE over the study period. Tillage system NUE was generally significantly higher than all other systems due to the mainly non-livestock nature of this system.

Figure 10-15: Nitrogen Use Efficiency: 3 year rolling average 2017-2022





Figure 10-16 illustrates that, on a three year rolling average basis across all farm systems, PUE (P outputs / P inputs) has generally declined from the start to the middle of the period and then increased towards the end of the period. However, farm gate PUE measures must be interpreted with care, since establishing true PUE at the individual far level requires a soil test.

Figure 10-16: Phosphorus Use Efficiency: 3 year rolling average 2017-2022



10.3 Social Sustainability Indicators

Figure 10-17 shows that on a three-year rolling average basis the rate of vulnerability (non-viable farm business and no off-farm employment) of all farming households has remained stable over the 2017-2021 period across all systems at between 33%-34% but declined to 28% at the end. Dairying and tillage systems tend to have significantly lower levels of household vulnerability than cattle and sheep systems.

Figure 10-17: Farm Household Vulnerability: 3 year rolling average 2017-2022




Figure 10-18 shows that on a three-year rolling average basis the percentage of farmers at risk of isolation was relatively static from the start to the end of the study period across all systems (except for sheep). However, overall isolation risk tends to be higher on non-dairy farms.

Figure 10-18: Isolation Risk: 3 year rolling average 2017-2022 (average per system)




Figure 10-19 shows that on a three year rolling average basis the percentage of all farms with a high age profile has increased between the start and end of the study period (25% to 35%). Dairy farms tend to have the lowest age profile across all the farm systems compared to other systems (15-17% towards the end of the study period) which tend to be double or treble this rate.

Figure 10-19: High Age Profile: 3 year rolling average 2017-2022 (average per system)




Figure 10-20 shows that the hours worked on-farm per annum has declined slightly across all farms systems except dairying between 2017 and 2022. Hours worked on farm per annum is significantly higher on dairy farms, compared to all other farm systems.

Figure 10-20: Hours Worked On Farm Per Annum: 3 year rolling average 2017-2022 (average per system)




Figure 10-21 illustrates the total hours worked by the farm operator (including on and off farm employment) by farm systems. Results indicate that longer overall hours are worked by dairy farmers and that the number of hours worked has increased over time. The opposite trend is observed for livestock farmers (fewer hours worked over time) with the number of hours worked by farmers on tillage farms increased slightly over the study period.

Figure 10-21: Total Hours Worked Per Annum: 3 year rolling average 2017-2022 (average per system)





Figure 10-22 indicates that the percentage of all farmers who have received some form of agricultural education has increased over the period 2017-2022, from 44% to 53%. Significantly, higher levels of formal agricultural education are observed for dairy farmers.

Figure 10-22: Formal Agricultural Education: 3 year rolling average 2017-2022 (average per system)




10.4 Environmental Emissions Intensity Trends

The following section examines the trends in environmental efficiency of production for the main products produced on livestock farms (milk, beef and sheep meat). Results are again reported on the basis of a three year rolling average (e.g. the 2015-2017 results are the average of 2015, 2016 and 2017 results). Results for individual years are reported in the appendices for each farm system.

Results presented in Figure 10-23 show that, the kg of CO₂ equivalent per kg of FPCM (IPCC based) has generally followed a declining trend since the middle of the period analysed.


Figure 10-23: Ag. GHG Emissions per kg FPCM: 2017-2022 Dairy Farms



Note: (IPCC approach) 3 year rolling average

Figure 10-24 indicates that kg of CO₂ equivalent per kg of live-weight beef produced on cattle farms also tended to follow a declining trend towards the end of the period analysed.


Figure 10-24: Ag. GHG Emissions per kg live-weight beef produced: 2017-2022 (Cattle Farms)



Note: (IPCC approach) 3 year rolling average

Figure 10-25 indicates, on a three year rolling average basis, a steady declining trend in terms of kg of CO₂ emitted per kg of live-weight sheep over the study period, despite an increase in the middle of the study period.


Figure 10-25: Ag. GHG Emissions per kg live-weight sheep produced: 2017-2022 Sheep Farms



Note: (IPCC approach) 3 year rolling average

Energy based GHG emissions may be affected by the weather in any given year (e.g. wet conditions may require extra movement of farm livestock herds). Results presented in Figure 10-26 indicate a gradual decline in energy based GHG emissions associated with milk production over the study period


Figure 10-26: Energy use related GHG emissions per kg FPCM: 2017-2022 Dairy Farms



Note: (IPCC approach) 3 year rolling average

Energy based CO₂ emissions related to the production of live-weight beef on cattle farms were relatively static over the study, with a declining trend evident toward the end of the study period as illustrated in Figure 10-27.


Figure 10-27: Energy use related GHG emissions per kg live-weight beef produced: 2017-2022 Cattle Farms



Note: (IPCC approach) 3 year rolling average

Energy related GHG emissions from the production of live-weight sheep tended to follow a declining trends over the course of the study period as illustrated in Figure 10-28.


Figure 10-28: Energy use related GHG emissions per kg live-weight sheep produced: 2017-2022 Sheep Farms



Note: (IPCC approach) 3 year rolling average


On a three year rolling average basis, the NH₃ emissions intensity of milk production tended to be decline towards the end of the study period as outlined in Figure 10-29.

Figure 10-29: Ammonia emissions per kg FPCM: 2017-20223 year rolling average Dairy Farms




On a three-year rolling average basis, NH₃ emissions per kg of live-weight beef produced on cattle farms also declined towards the end of the study as shown in Figure 10-30.

Figure 10-30: Ammonia emissions per kg live-weight beef produced: 2017-2022 Cattle Farms



The NH₃ emissions per kg of live-weight sheep meat produced on sheep farms was seen to decline towards the end of the study period, as illustrated in Figure 10-31.

Figure 10-31: Ammonia emissions per kg live-weight sheep produced: 2017-2022 Sheep Farms



References

- Breen, J.P., Donnellan, T., Westhoff, P., 2010. Food for Thought: EU Climate Change Policy Presents New Challenges for Agriculture. EuroChoices 9, 24-29.
- Buckley, C., Wall, D.P., Moran, B., Murphy, P.N.C., 2015. Developing the EU Farm Accountancy Data Network to derive indicators around the sustainable use of nitrogen and phosphorus at farm level. Nutrient Cycling in Agroecosystems 102, 319-333.
- Buckley, C., Donnellan, T., Dillon, E., Hanrahan, K., Moran, B., Ryan, M., 2019. Teagasc National Farm Survey 2017 Sustainability Report. Available: <https://www.teagasc.ie/media/website/publications/2019/2017-sustainability-report-250319.pdf>
- Buckley, C and Donnellan, T., 2020a. Teagasc National Farm Survey 2018 Sustainability Report. Available: <https://www.teagasc.ie/media/website/publications/2020/Teagasc-NFS-2018-Sustainability-Report.pdf>
- Buckley, C and Donnellan, T., 2020b. Teagasc National Farm Survey 2019 Sustainability Report. Available: <https://www.teagasc.ie/media/website/publications/2020/NFS-2019-Sustainability-Report.pdf>
- Buckley, C and Donnellan, T., 2021. Teagasc National Farm Survey 2020 Sustainability Report. Available: <https://www.teagasc.ie/media/website/publications/2021/2020-Sustainability-Report.pdf>
- Buckley, C and Donnellan, T., 2022. Teagasc National Farm Survey 2021 Sustainability Report. Available: <https://www.teagasc.ie/media/website/publications/2022/2021-Sustainability-Report.pdf>
- Central Statistics Office, 2022. Ireland: Land Use Categories 1990-2020. Available: www.cso.ie/en/releasesandpublications/ep/p-eii/environmentalindicatorsireland2022/landuse/
- Central Statistics Office, 2023a. National Average Price (Euro) for selected consumer items. Available: <https://data.cso.ie/#>
- Central Statistics Office, 2023b. Agricultural Output and Input price Indices. Available: <https://data.cso.ie/table/AHA04>
- Dillon, E., Hennessy, T., Moran, B., Lennon, J., Lynch, J., Brennan, M. and Donnellan, T., 2017. Small Farms 2015 - Teagasc National Farm Survey. Available: <https://www.teagasc.ie/media/website/publications/2017/Small-Farms-Survey.pdf>
- Donnellan, T., Moran, B., Lennon, J., & Dillon, E., 2020. Teagasc National Farm Survey 2019 Preliminary Results. Available: <https://www.teagasc.ie/media/website/publications/2020/TeagascNFS2019-Preliminary-Results.pdf>
- Duffy, P., Black, K., Fahey, D., Hyde, B., Kehoe, A., Murphy, J., Quirke, B., Ryan, A.M. & Ponzi, J. 2023. Ireland National inventory report 2022 greenhouse gas emissions 1990 - 2020 reported to the United Nations framework convention on climate change. Available at: https://www.epa.ie/publications/monitoring--assessment/climate-change/air-emissions/Ireland-NIR-2022_Merge_v2..pdf
- Duffy, P., Hyde, B., Ryan, A.M. Murphy, J., Quirke, B., Fahey, D., & Kehoe, A., 2020b. Informative inventory report 2020 air pollutant emissions in Ireland 1990–2018 reported to the secretariat

of the unece convention on long-range transboundary air pollution and to the European Union. Available at:

https://www.epa.ie/pubs/reports/air/airemissions/airpollutantemissions/iir2019/IIR_2019_Final%2001.05.19_Cover.v2.pdf

EPA. 2022. Ireland's Air Pollutant Emissions 1990-2030. Environmental Protection Agency. Available at: https://www.epa.ie/publications/monitoring--assessment/climate-change/air-emissions/Air-Pollutant-Emissions-Report-2022_final.pdf

EPA, 2023. Environmental Indicators. Climate – Greenhouse gas by sector in 2020. Available at: <https://www.epa.ie/our-services/monitoring--assessment/climate-change/ghg/latest-emissions-data/>

European Commission, 2016. Directive 2016/2284 of the European Parliament and of the Council of 14 December 2016 on the reduction of national emissions of certain atmospheric pollutants, amending Directive 2003/35/EC and repealing Directive 2001/81/EC Available at: <https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32016L2284&from=EN> [accessed 20/03/2020] .

European Commission, 2020. A farm to fork Strategy, for a fair, healthy and environmentally friendly food system. Available: https://ec.europa.eu/food/farm2fork_en

Government of Ireland, 2021a. National Minimum Wage. Available: <https://www.gov.ie/en/publication/41a981-the-minimum-wage-in-ireland/>

Government of Ireland, 2020a, The history of the minimum wage. Available: <https://www.gov.ie/en/publication/9463f6-historic-nmw-rates/#>

Government of Ireland, 2021b. Climate Action and Low Carbon Development (Amendment) Act 2021 Available: <https://data.oireachtas.ie/ie/oireachtas/act/2021/32/eng/enacted/a3221.pdf>

Government of Ireland, 2022. Government announces sectoral emissions ceilings, setting Ireland on a pathway to turn the tide on climate change. Available: <https://www.gov.ie/en/press-release/dab6d-government-announces-sectoral-emissions-ceilings-setting-ireland-on-a-pathway-to-turn-the-tide-on-climate-change/>

Hennessy, T., Buckley, C., Dillon, E., Donnellan, T., Hanrahan, K., Moran, B., Ryan, M., 2013. Assessing the Sustainability of Irish Farms - Teagasc National Farm Survey. Agricultural Economics & Farm Surveys Department, REDP, Teagasc, Athenry, Co. Galway. Available: <https://www.teagasc.ie/media/website/publications/2013/SustainabilityReport.pdf>

Herron, J., Hennessy, D., Curran, T., Moloney, A, O'Brien, D., 2021. Journal of Dairy Science Vol. 104 No. 1, pp. 7902-7918.

Murphy, E., 2022. Sustainability Data & Analytics Manager, Bord Bia. Personal Communication.

Lynch, J., Hennessy, T., Buckley, C., Dillon, E., Donnellan, T., Hanrahan, K., Moran, B., Ryan, M., 2016. Teagasc National Farm Survey 2015 Sustainability Report. Teagasc National Farm Survey. Agricultural Economics & Farm Surveys Department, REDP, Teagasc, Athenry, Co. Galway. ISBN 978-1-84170-631-3. Available: <https://www.teagasc.ie/media/website/publications/2017/2015-sustainability-report.pdf>

O'Brien, D., Capper, J.L., Garnsworthy, P.C., Grainger, C., Shalloo, L., 2014. A case study of the carbon footprint of milk from high-performing confinement and grass-based dairy farms. *Journal of Dairy Science*, 97, 1835-1851.

Sustainability Energy Authority of Ireland, 2021. Commercial Fuel Cost Archive. Available:

<https://www.seai.ie/resources/publications/Commercial-Fuel-Cost-Archives.pdf>

Sutton, M.A., Moncrieff, J.B. & Fowler, D., 1992. Deposition of atmospheric ammonia to moorlands. *Environmental Pollution*, 75, 15–24.

Teagasc, 2022. Soil Results. Available: <https://www.teagasc.ie/media/website/crops/soil-and-soil-fertility/Teagasc-Soil-Fertility-Report-2022.pdf>

World Bank (2020). Agricultural Land (% of land area) Ireland. Available:

<https://data.worldbank.org/indicator/AG.LND.AGRI.ZS>

Appendix 1

01 Individual year results by farm system 2017-2022



Appendix 1 – Individual year results by farm system 2017-2022

Table A 1: Sustainability Indicator results for Dairying Farms 2017-2022

<i>Indicator</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>	<i>2021</i>	<i>2022</i>
Economic Sustainability Metrics						
Economic return per hectare (gross output)	3,722	3,641	3,620	3,730	4,324	6,005
Profitability per hectare (gross margin)	2,113	1,730	1,802	1,920	2,396	3,509
Family farm income per hectare	1,537	1,059	1,122	1,246	1,548	2,332
Productivity of labour	69,449	48,986	52,449	57,813	73,941	109,003
Market orientation	90%	89%	88%	90%	91%	94%
Viability	85%	73%	75%	80%	86%	93%
Social Sustainability Metrics						
Household vulnerable	8%	13%	12%	8%	7%	4%
Isolation	8%	7%	7%	6%	6%	6%
High age profile	12%	12%	15%	16%	16%	18%
Hours worked on farm	2,345	2,398	2,380	2,422	2,545	2,498
Total hours worked	2,487	2,552	2,545	2,586	2,725	2,657
Agricultural education	76%	77%	79%	83%	82%	83%
Environmental Sustainability Metrics						
	tonnes CO ₂ eqv per farm					
Total farm average Ag. GHG emissions	547.4	575.2	579.0	600.4	616.1	606.0
of which dairy	375.6	393.8	410.7	431.1	448.5	443.3
cattle	170.2	180.2	167.3	168.6	166.4	161.2
sheep	1.5	1.2	1.0	0.7	1.2	1.5
other	0.0	0.0	0.0	0.0	0.0	0.0
energy use	17.9	18.3	18.4	17.8	20.5	19.7
	tonnes CO ₂ eqv per ha					
Ag GHG Emissions	9.3	9.5	9.4	9.4	9.6	9.4
Energy GHG Emissions	0.33	0.33	0.32	0.30	0.35	0.33
kg CO ₂ eqv						
Ag. GHG Emissions per kg milk	0.89	0.90	0.88	0.87	0.85	0.84
Ag. GHG Emissions per kg FPCM	0.95	0.91	0.88	0.85	0.84	0.82
Ag. GHG Emissions per € output	2.9	3.1	2.8	2.9	2.5	1.7
Energy GHG Emissions per kg milk	0.053	0.046	0.044	0.038	0.037	0.035
Energy GHG Emissions per kg FPCM	0.037	0.037	0.034	0.032	0.036	0.033
Energy GHG Emissions per € output	0.107	0.109	0.098	0.095	0.094	0.061
GHG Emissions per kg FPCM (LCA)	1.12	1.12	1.07	1.07	1.05	1.06
	tonnes NH ₃ per farm					
Total farm average NH ₃ emissions	2.82	3.09	3.03	3.07	2.89	3.00
of which dairy	1.94	2.15	2.18	2.23	2.13	2.24
cattle	0.87	0.93	0.85	0.83	0.76	0.76
sheep	0.01	0.01	0.01	0.01	0.00	0.00
tillage	0.00	0.00	0.00	0.00	0.00	0.00

Teagasc National Farm Survey 2022 Sustainability Report

Indicator	2017	2018	2019	2020	2021	2022
Kg NH ₃						
NH3 emissions per hectare	48.0	51.4	49.1	48.2	45.1	46.9
NH3 emissions per Euro output	0.015	0.017	0.014	0.015	0.012	0.009
NH3 emissions per kg milk	0.005	0.005	0.005	0.004	0.004	0.004
NH3 emissions per kg FPCM	0.005	0.005	0.005	0.004	0.004	0.004
N Balance per hectare	172.3	201.0	179.9	175.6	163.7	158.6
P Balance per hectare	11.4	15.7	13.3	12.4	12.9	10.4
percentage						
N use efficiency	24.3	21.5	24.2	25.5	26.8	27.7
P use efficiency	58.3	47.7	53.3	55.4	53.3	60.8
Innovation Metrics						
Discussion Group Membership	43%	42%	44%	44%	45%	41%
Milk Recording	41%	38%	46%	43%	48%	50%
% of slurry spread using LESS	5%	5%	32%	52%	67%	75%
% of slurry applied during spring	56%	51%	54%	53%	51%	56%
% chemical N applied as Protected Urea	0%	0%	3%	5%	7%	14%
% of farms reseeded	28%	26%	25%	32%	31%	30%
% of farms liming	33%	30%	36%	38%	44%	53%

Table A 2: Sustainability Indicator results for Cattle Farms 2017-2022

<i>Indicator</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>	<i>2021</i>	<i>2022</i>
Economic Sustainability Metrics						
Economic return per hectare (gross output)	1,338	1,316	1,309	1,324	1,532	1,740
Profitability per hectare (gross margin)	536	486	495	511	653	845
Family Farm Income per hectare	393	321	310	333	404	372
Productivity of labour	17,035	13,510	13,831	14,255	17,445	17,154
Market orientation	64%	62%	60%	62%	66%	70%
Viability	25%	18%	19%	18%	27%	27%
Social Sustainability Metrics						
Household vulnerable	39%	39%	41%	39%	34%	33%
Isolation	22%	24%	23%	23%	18%	17%
High age profile	32%	38%	39%	40%	40%	41%
Hours worked	1,510	1,521	1,480	1,494	1,486	1,370
Total Hours Worked	2,203	2,215	2,161	2,173	2,221	2,139
Agricultural education	38%	40%	40%	43%	46%	49%
Environmental Sustainability Metrics						
tonnes CO ₂ eqv per farm						
Total farm average Ag. GHG emissions	152.7	161.6	150.4	147.8	156.9	150.9
of which dairy	0.0	0.0	0.0	0.0	0.0	0.0
cattle	148.1	156.6	145.6	142.8	152.0	146.6
sheep	4.3	4.7	4.5	4.8	4.5	3.9
other	0.3	0.2	0.2	0.2	0.5	0.4
energy use	6.2	6.1	5.9	5.6	6.1	5.8
tonnes CO ₂ eqv per ha						
Ag GHG Emissions	4.7	4.8	4.5	4.4	4.7	4.4
Energy GHG Emissions	0.2	0.2	0.2	0.2	0.2	0.2
kg CO ₂ eqv						
Ag. GHG Emissions per kg live-weight beef	12.1	12.7	11.8	11.8	12.1	9.4
Ag. GHG Emissions per € output	6.5	5.9	5.4	5.1	4.5	3.6
Energy GHG Emissions per kg live-weight beef	0.53	0.52	0.49	0.48	0.51	0.39
Energy GHG Emissions per € output	0.26	0.25	0.22	0.21	0.19	0.15
tonnes NH ₃ per farm						
Total farm average NH ₃ emissions	0.84	0.89	0.82	0.76	0.79	0.76
of which dairy	0.00	0.00	0.00	0.00	0.00	0.00
cattle	0.82	0.87	0.80	0.74	0.77	0.75
sheep	0.02	0.02	0.02	0.02	0.02	0.01
tillage	0.00	0.00	0.00	0.00	0.00	0.00
Kg NH ₃						
NH ₃ emissions per hectare	26.1	26.2	24.3	22.9	23.1	22.0
NH ₃ emissions per Euro output	0.04	0.03	0.03	0.03	0.02	0.02
NH ₃ emissions per kg live-weight beef	0.07	0.07	0.07	0.06	0.07	0.05
kg per ha						
N Balance per hectare	58.9	67.2	57.7	58.9	63.3	43.6
P Balance per hectare	5.2	5.7	4.9	4.8	6.2	2.4

Teagasc National Farm Survey 2022 Sustainability Report

Indicator	2017	2018	2019	2020	2021	2022
			percentage			
N use efficiency	22.8	20.6	22.3	23.0	21.9	33.9
P use efficiency	66.9	61.1	67.6	68.9	63.3	89.2
			Per kg of N Surplus			
Innovation Metrics						
Discussion Group Membership	17%	21%	19%	11%	16%	14%
% of slurry spread using LESS	2%	3%	12%	18%	25%	34%
% of slurry applied during spring	48%	43%	41%	44%	49%	44%
% chemical N applied as Protected Urea	0%	0%	1%	2%	2%	4%
% of farms reseeded	6%	9%	8%	9%	14%	11%
% of farms liming	15%	14%	11%	11%	21%	25%

Table A 3: Sustainability Indicator results for Sheep Farms 2017-2022

<i>Indicator</i>	2017	2018	2019	2020	2021	2022
Economic Sustainability Metrics						
Economic return per hectare (gross output)	1,382	1,314	1,241	1,293	1,476	1,475
Profitability per hectare (gross margin)	549	396	409	489	623	625
Family Farm Income per hectare	466	275	321	401	445	340
Productivity of labour	17,224	12,363	14,256	17,652	18,725	14,890
Market orientation	60%	59%	55%	61%	66%	65%
Viability	28%	20%	23%	25%	39%	26%
Social Sustainability Metrics						
Household vulnerable	41%	45%	41%	39%	27%	29%
Isolation	12%	13%	23%	26%	21%	13%
High age profile	30%	38%	40%	46%	37%	28%
Hours worked on farm	1,644	1,579	1,555	1,524	1,541	1,477
Total hours worked	2,132	1,996	2,055	1,988	2,085	2,246
Agricultural education	42%	50%	53%	52%	57%	61%
Environmental Sustainability Metrics						
tonnes CO ₂ eqv per farm						
Total farm average Ag. GHG emissions	162.6	159.6	153.9	154.0	165.9	136.2
of which dairy	1.8	0.0	0.0	0.0	0.0	0.5
cattle	77.2	76.5	72.4	72.0	75.3	61.2
sheep	83.3	82.8	81.3	81.6	90.1	74.0
other	0.3	0.3	0.2	0.4	0.6	0.4
energy use	6.5	6.2	6.1	5.3	6.0	5.5
tonnes CO ₂ eqv per ha						
Ag GHG Emissions	4.3	4.1	3.9	3.8	4.1	3.3
Energy GHG Emissions	0.20	0.17	0.17	0.14	0.17	0.16
kg CO ₂ eqv						
Ag. GHG Emissions per kg live-weight sheep produced	11.2	12.1	10.8	12.2	11.5	7.8
Ag. GHG Emissions per € output	5.1	5.0	5.6	4.5	4.0	3.5
Energy Emissions per kg live-weight sheep produced	0.55	0.52	0.5	0.49	0.51	0.53
Energy GHG Emissions per € output	0.28	0.23	0.23	0.17	0.16	0.09
tonnes NH ₃ per farm						
Total farm average NH ₃ emissions	0.79	0.79	0.77	0.74	0.78	0.56
of which dairy	0.01	0.00	0.00	0.00	0.00	0.00
cattle	0.45	0.45	0.43	0.41	0.42	0.36
sheep	0.34	0.34	0.34	0.33	0.36	0.20
tillage	0.00	0.00	0.00	0.00	0.00	0.00
kg NH ₃						
NH ₃ emissions per hectare	20.7	20.4	19.2	18.6	19.4	12.7
NH ₃ emissions per Euro output	0.02	0.03	0.03	0.02	0.02	0.01
NH ₃ emissions per kg live-weight sheep	0.04	0.05	0.04	0.05	0.04	0.02
kg per ha						
N Balance per hectare	50.7	64.1	47.0	49.0	53.4	30.4
P Balance per hectare	5.8	7.0	4.8	5.8	7.1	3.3

Teagasc National Farm Survey 2022 Sustainability Report

<i>Indicator</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>	<i>2021</i>	<i>2022</i>
	percentage					
N use efficiency	30.6	24.6	29.4	29.3	25.7	37.6
P use efficiency	62.1	52.2	60.8	58.6	57.8	78.3
Innovation Metrics						
Discussion Group Membership	28%	26%	25%	16%	19%	17%
% of slurry applied during spring (where slurry is generated)	26%	22%	28%	31%	33%	20%
% chemical N applied as Protected Urea	0%	0%	0%	2%	2%	2%
% of farms reseeded	3%	16%	7%	8%	13%	12%
% of farms liming	22%	17%	12%	16%	28%	24%

Table A 4: Sustainability Indicator results for Tillage Farms 2017-2022

<i>Indicator</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>	<i>2021</i>	<i>2022</i>
Economic Sustainability Metrics						
Economic return per hectare (gross output)	1,737	1,855	1,966	1,693	2,218	2,812
Profitability per hectare (gross margin)	819	902	878	753	1,189	1,581
Family farm income per hectare	618	656	584	550	839	1,043
Productivity of labour	44,591	43,928	36,684	38,537	75,185	97,910
Market orientation	76%	79%	75%	73%	79%	82%
Viability	74%	63%	62%	66%	76%	79%
Social Sustainability Metrics						
Household vulnerable	11%	17%	17%	17%	13%	10%
Isolation	19%	16%	17%	16%	19%	18%
High age profile	27%	33%	32%	37%	28%	25%
Hours worked on farm	1,466	1,510	1,539	1,422	1,519	1,416
Total hours worked	1,937	2,081	2,220	2,062	2,215	2,133
Agricultural education	61%	54%	64%	56%	63%	70%
Environmental Sustainability Metrics						
tonnes CO ₂ eqv per farm						
Total farm average Ag. GHG emissions	133.9	146.2	144.2	146.6	152.3	137.5
of which dairy	0.0	0.0	0.0	0.0	0.0	0.0
cattle	98.2	111.2	103.4	109.7	103.8	85.6
sheep	9.4	10.2	12.6	15.1	15.1	12.9
other	26.4	24.7	28.2	21.8	33.4	40.0
energy use	13.6	14.4	14.2	13.0	16.5	12.8
tonnes CO ₂ eqv per ha						
Ag GHG Emissions	2.1	2.4	2.2	2.2	2.2	1.9
Energy GHG Emissions	0.23	0.22	0.23	0.19	0.22	0.21
kg CO ₂ eqv						
Ag. GHG Emissions per € output	1.66	1.65	1.66	1.78	1.31	0.86
Energy GHG Emissions per € output	0.16	0.15	0.16	0.14	0.12	0.12
tonnes NH ₃ per farm						
Total farm average NH ₃ emissions	0.60	0.74	0.74	0.68	0.73	0.72
of which dairy	0.00	0.00	0.00	0.00	0.00	0.00
cattle	0.51	0.57	0.51	0.54	0.49	0.42
sheep	0.04	0.04	0.05	0.06	0.05	0.03
tillage	0.05	0.13	0.16	0.08	0.19	0.27
kg NH ₃						
NH ₃ emissions per hectare	9.0	10.8	10.0	10.1	9.8	9.1
NH ₃ emissions per Euro output	0.01	0.01	0.01	0.01	0.01	0.01
kg per ha						
N Balance per hectare	41.4	61.9	45.8	44.2	39.4	40.5
P Balance per hectare	4.4	9.9	6.1	9.2	6.5	3.6
percentage						
N use efficiency	69.7	56.8	65.6	61.8	65.9	66.6
P use efficiency	86.4	70.4	82.2	70.5	84.8	90.4

Teagasc National Farm Survey 2022 Sustainability Report

<i>Indicator</i>	<i>2017</i>	<i>2018</i>	<i>2019</i>	<i>2020</i>	<i>2021</i>	<i>2022</i>
Innovation Metrics						
Discussion Group Membership	24%	21%	29%	21%	26%	30%
Break Crop	9%	19%	19%	15%	12%	15%
% of farms liming	32%	27%	28%	20%	32%	26%

Table A 5: Sustainability Indicator results for All Farms 2017-2022

<i>Indicator</i>	2017	2018	2019	2020	2021	2022
Economic Sustainability Metrics						
Economic return per hectare (gross output)	1,800	1,774	1,766	1,787	2,085	2,556
Profitability per hectare (gross margin)	840	727	746	783	1,006	1,307
Family Farm income per hectare	626	471	479	528	652	775
Productivity of labour	28,506	21,941	22,544	24,588	32,271	39,571
Market orientation	69%	68%	65%	68%	72%	75%
Viability	40%	32%	33%	34%	44%	43%
Social Sustainability Metrics						
Household vulnerable	31%	34%	34%	32%	26%	25%
Isolation	18%	18%	19%	20%	17%	15%
High age profile	28%	33%	35%	36%	34%	34%
Hours worked on farm	1,678	1,688	1,660	1,663	1,691	1,597
Total hours worked	2,222	2,230	2,218	2,209	2,291	2,251
Agricultural education	47%	50%	51%	53%	56%	59%
Environmental Sustainability Metrics						
	tonnes CO ₂ eqv per farm					
Total farm average Ag. GHG emissions	223.2	234.5	228.2	231.5	242.1	231.7
energy use	8.9	8.9	8.8	8.4	9.5	8.9
	tonnes CO ₂ eqv					
Ag GHG Emissions per hectare	5.3	5.4	5.1	5.1	5.3	5.0
Ag GHG Emissions per Euro output	5.3	4.9	4.7	4.4	3.8	3.1
Energy GHG Emissions per hectare	0.23	0.22	0.21	0.20	0.22	0.21
Energy Emissions per Euro output	0.23	0.21	0.20	0.18	0.16	0.19
	tonnes NH ₃ per farm					
Total farm average NH ₃ emissions	1.17	1.26	1.21	1.17	1.17	1.13
	kg NH ₃					
NH ₃ emissions per hectare	27.8	28.6	26.9	25.9	25.5	24.0
NH ₃ emissions per Euro output	0.03	0.03	0.02	0.02	0.02	0.01
	kg per ha					
N Balance per hectare	74.4	88.8	75.3	76.1	78.2	60.4
P Balance per hectare	6.1	7.7	6.2	6.4	7.4	3.9
	percentage					
N use efficiency	27.2	24.0	26.6	27.0	25.5	35.1
P use efficiency	66.5	58.4	65.4	65.2	62.9	82.7

